

ON THE LATTICE OF SUBGROUPS OF THE LAMPLIGHTER GROUP

R. GRIGORCHUK AND R. KRAVCHENKO

ABSTRACT. The paper is devoted to the study of the lattice of subgroups of the Lamplighter type groups and to the relative gradient rank.

1. INTRODUCTION

The group \mathcal{L} called Lamplighter is defined as the wreath product $(\mathbb{Z}/2)\wr\mathbb{Z}$. This group and its generalizations like wreath products of two groups and, more generally, iterated wreath products play important role not only in the group theory but also in theory of random walks [23], [11], [5], and other areas of mathematics. One of the most unusual properties of \mathcal{L} is that the spectrum of Laplace operator on Cayley graph of \mathcal{L} (constructed using a special system of generators) is pure point spectrum and accordingly the spectral measure is discrete [19]. This result together with some other results which also involve the lamplighter type groups lead to the solution in ([18], [4], [12], [26], [29]) of different versions of the Atiyah Problem [3].

The group \mathcal{L} has a presentation

$$\langle a, b | b^2, [b, a^j b a^{-j}] \text{ for } j \geq 1 \rangle,$$

which is minimal (hence the group is not finitely presented). \mathcal{L} is a metabelian group of exponential growth and is a self-similar group generated by 2-state automaton over binary alphabet, [19], [17], [6]. Despite relatively simple algebraic structure, the group \mathcal{L} in fact is not so simple to study and there are many open questions related to it.

In this note we study a subgroup structure of \mathcal{L} and of groups $\mathcal{L}_{n,p} = (\mathbb{Z}/p\mathbb{Z})^n \wr \mathbb{Z}$, where p is a prime. Groups $\mathcal{L}_{1,p}$ play especially important role in study of solvable groups. For instance the result of Kropholler ([24]) tells that the groups of finite rank in the class of finitely generated solvable groups are characterized as those groups which have no section isomorphic with $\mathcal{L}_{1,p}$ for some p .

The first author was supported by Simons Foundation and Mittag-Leffler Institute. Both authors were supported by the ERC starting grant GA 257110 RaWG.

In section 3 we realize $\mathcal{L}_{1,p}$ as a group generated by finite automaton of Mealy type with p states over an alphabet on p symbols. This allows to realize $\mathcal{L}_{1,p}$ as a group acting on a p -ry rooted tree in a self-similar manner. Also this allows to identify a big part of the lattice of p -power subgroups with a set of vertex stabilizers for the action of a group on a tree. Our goal (not achieved yet) is complete understanding of the lattice of subgroups of $\mathcal{L}_{n,p}$. Even for subgroups of finite index or normal subgroups this is not an easy problem. Our first result is the description up to isomorphism of all subgroups of $\mathcal{L}_{n,p}$. In what follows we will skip index p , keeping it fixed while n can vary, and denote the group just by \mathcal{L}_n , to lighten the notation. We show in theorem 3.5 that either a subgroup of \mathcal{L}_n sits inside the base group \mathcal{A}_n of wreath product and hence is abelian, or it is isomorphic to \mathcal{L}_k for some k . As a corollary of our arguments we get that a subgroup of index p of \mathcal{L}_n is isomorphic to \mathcal{L}_n or to \mathcal{L}_{pn} (both cases occur). This will show in particular that \mathcal{L}_n has a descending chain with trivial intersection of subgroups of primary index a power of p that consists of groups isomorphic to \mathcal{L}_n . Groups with this property are called scale invariant. Around 2005 there was a conjecture supported by Benjamini and some other mathematicians that finitely generated scale invariant groups are virtually nilpotent. The conjecture is not correct as we see. That \mathcal{L} is scale invariant was implicitly established in [19] and in details investigated by V. Nekrasevych and Gábor Pete in [28], where many other scale invariant groups are constructed.

In Theorem 3.11 we describe all maximal subgroups of \mathcal{L}_n (they happen to be all of finite index).

Recall that a subgroup H of a group G is called weakly maximal (or near maximal, see [31]) if it has infinite index, but every subgroup of G containing H as a proper subgroup has finite index in G . All weakly maximal subgroups of \mathcal{L}_n are described in Theorem 3.12, moreover it is determined which of them are closed in pro- p topology.

A property of \mathcal{L}_n called *hereditary free from finite normal subgroups* is established which, according to results of [15], allows to conclude that any action of \mathcal{L}_n on the rooted tree induces topologically free action on the boundary of the tree. We get a formulae for a number of subgroups of \mathcal{L}_n of index m (Proposition 3.9) which allows to obtain an upper bound on subgroup growth (Proposition 3.10). A description of pro- p -completion of \mathcal{L}_n is presented in Lemma 3.30.

Our main tool to study subgroup structure of \mathcal{L}_n is based on technique of association to a group a suitable module on which group acts. For the case of solvable groups this technique was successfully used by

P.Hall and many other researchers [21]. We also follow this line considering the base subgroup as a module over the ring of Laurent polynomial in one variable (in fact different module structures are used). This allows us to associate to a subgroup $H \leq \mathcal{L}_n$ a triple (s, V_0, v) consisting of $s \in \mathbb{N}$, $V_0 \leq \mathcal{A}_n$, $v \in \mathcal{A}_n$ that satisfies certain properties listed in Lemma 3.1 and Corollary 3.4. In case of a normal subgroup the triple satisfies more constraints (Lemmas 3.2 and 3.20). Theorem 3.22 gives a description of the lattice of normal subgroups of $\mathcal{L}_{1,p}$. As a result we make some contribution to the normal subgroup growth (Theorem 3.17 and Corollary 3.19). Theorems 3.25 and 3.35 describe when the subgroup is closed in profinite and pro- p topologies in terms of the associated triples.

The last part of the paper deals with one interesting recent notion of rank gradient of a subgroup. It was introduced by Lackenby and studied by him, Abert, Jaikin-Zapirain, Luck, Nikolov, and other researchers. In many cases (in particular under certain amenability assumptions) it is equal to zero. A basic open question about gradient rank in the situation when a group is amenable is Question 1 formulated at section 4.

In a situation of zero gradient rank and when a subgroup $H \leq G$ is presented as the intersection of descending sequence of subgroups $\{H_m\}$ of finite index it is reasonable to study a relation between the index $[G : H_m]$ and rank $rk(H_m)$ (i.e. a minimal number of generators) as a function $rg(m)$ of natural argument m . It is interesting to know what type of asymptotic of such functions arises, how this function, which we call relative gradient rank, depends on presentation of subgroup in the form of intersection etc. Very little is known about this. The first observation in this direction in the case of lamplighter group was made in [2], Theorem 4.2 provides further information.

After the case of Lamplighter type groups is treated the next step would be to consider the case of metabelian groups and more generally of solvable groups and even of elementary amenable groups [9], [16].

Another option is the study of the case of nonelementary amenable groups starting with the 3-generated 2-group of intermediate growth constructed by the first author in [20].

Acknowledgements. We would like to express our thanks to Y. de Cornulier and D.Segal. We are also indebted to Anna Erschler and Laboratoire de Mathématiques d'Orsay for their hospitality. The major part of the work was done while the first author was visiting it and the second author was a postdoc there.

2. PRELIMINARY FACTS

We consider groups

$$\mathcal{L}_n = (\mathbb{Z}/p\mathbb{Z})^n \wr \mathbb{Z} = \oplus_{\mathbb{Z}} (\mathbb{Z}/p\mathbb{Z})^n \rtimes \mathbb{Z}.$$

Let \mathcal{A}_n denote the subgroup $\oplus_{\mathbb{Z}} (\mathbb{Z}/p\mathbb{Z})^n$ of \mathcal{L}_n . Let x denote the right shift by one on \mathcal{A}_n , considered as a generator of active group \mathbb{Z} . Elements of \mathcal{L}_n are written as (v, s) where $v \in \mathcal{A}_n$ and $s \in \mathbb{Z}$. If $g = (v, s)$ and $h = (w, t)$ are elements of \mathcal{L}_n then the direct computation gives that $gh = (v + x^s w, s + t)$ and $g^{-1} = (-x^{-s} v, -s)$.

2.1. The ring $R = \mathbb{F}_p[x, x^{-1}]$. Let \mathbb{F}_p be the simple field of characteristic p . In the following we will make repeated use of the properties of the ring $\mathbb{F}_p[x, x^{-1}]$ of Laurent polynomials, which we denote by R . Note also that R is just $\mathbb{F}_p[\mathbb{Z}]$, the group ring of \mathbb{Z} over \mathbb{F}_p .

Lemma 2.1. *R is principal ideal domain. The invertible elements of R are $\{ax^q | a \in \mathbb{F}_p^*, q \in \mathbb{Z}\}$. Each ideal I of R is of the form Rf where $f \in \mathbb{F}_p[x]$, $f(0) = 1$, such f is uniquely defined by I , up to multiplication by a nonzero element of \mathbb{F}_p . The factor ring R/I is isomorphic to $\mathbb{F}_p[x]/f\mathbb{F}_p[x]$. Therefore, each element of R/I has the unique representative polynomial g with degree of g less than f .*

Proof. Let I be an ideal in R . Note that $\mathbb{F}_p[x] \subset R$. Then $I \cap \mathbb{F}_p[x]$ is an ideal in $\mathbb{F}_p[x]$ and since it is principal ideal domain $I \cap \mathbb{F}_p[x] = f\mathbb{F}_p[x]$ for some $f \in \mathbb{F}_p[x]$. Then $Rf \subset I$. For $h \in I$ $h = x^{-k}g$ for some $k \in \mathbb{N}$ and $g \in \mathbb{F}_p[x]$. Thus $g \in I \cap \mathbb{F}_p[x] = f\mathbb{F}_p[x]$, and so $h = x^{-k}fa \in Rf$. Therefore $Rf = I$. Note that if $f = x^k f_0$ and $f_0(0) = 1$ then $Rf = Rf_0$. Thus we have proved that each ideal of R is generated by a single element of the form $f \in \mathbb{F}_p[x]$, $f(0) = 1$. If $Rf_1 = Rf_2$ then $f_1 = uf_2$ with $u \in R$ invertible in R , therefore $u = ax^k$. But since both $f_i(0) = 1$ we have that $k = 0$.

To prove that R/I is isomorphic to $\mathbb{F}_p[x]/f\mathbb{F}_p[x]$ when $I = Rf$, note that $\mathbb{F}_p[x]$ is a subring of R and $I \cap \mathbb{F}_p[x] = f\mathbb{F}_p[x]$, therefore $\mathbb{F}_p[x]/f\mathbb{F}_p[x]$ is subring of R/I . It is left to note that since $f(0) = 1$ we have that $f = a + xg$ for some $g \in \mathbb{F}_p[x]$ and $a \in \mathbb{F}_p^*$ and thus x is invertible in $\mathbb{F}_p[x]/f\mathbb{F}_p[x]$. \square

Note that it follows in particular that for $f, f_1 \in \mathbb{F}_p[x]$ with $f(0) \neq 0$, f_1 divides f in R if and only if f_1 divides f in $\mathbb{F}_p[x]$. Therefore in the following we will just say that f_1 divides f and write $f_1 | f$ meaning that it happens both in R and in $\mathbb{F}_p[x]$.

Definition 2.2. *Let $f \in R$. $\text{Deg}(f)$ is the dimension $\dim_{\mathbb{F}_p} R/fR$ of vector space R/fR over \mathbb{F}_p . Note that it is also equal to the $m - k$,*

where m is the largest degree of x in f with the nonzero coefficient and k is the smallest degree of x with nonzero coefficient.

Note that we use capital 'D' here in order to avoid confusion with standard notion of degree when $f \in \mathbb{F}_p[x]$. In this case $\deg f$ is the dimension of $\mathbb{F}_p[x]/f\mathbb{F}_p[x]$, and $\text{Deg } f$ is the dimension of R/fR , clearly $\text{Deg } f \leq \deg f$ and they are equal if and only if $f(0) \neq 0$.

We will be also interested in the following facts about the ring $M_n(R)$ of $n \times n$ matrices over R and the right module R^n over $M_n(R)$ with the standard action of $M_n(R)$ (but often we also consider R^n as a left module over R with the action described in each particular case).

The next lemma is basic for many considerations (and true for matrices with coefficients in any principal ideal domain).

Lemma 2.3. *Let $g \in M_n(R)$. Then there exist $a, b \in M_n(R)$ such that $g = adb$ where $d \in M_n(\mathbb{F}_p[x])$ is the diagonal matrix, and $\det(a), \det(b)$ are invertible in R (i.e. $a, b \in GL_n(R)$).*

Corollary 2.4. $\dim_{\mathbb{F}_p} R^n/gR^n = \text{Deg}(\det(g))$.

Proof. Indeed, if $g = adb$ is the decomposition then $\det(g)$ equals to a product of $\det(d)$ and an invertible element of R , thus $R/\det(g)R = R/\det(d)R$ and therefore $\text{Deg}(\det(g)) = \text{Deg}(\det(d))$. Also, $R^n/gR^n = R/adR^n$ is isomorphic to R^n/dR^n . Now if d_1, \dots, d_n are diagonal entries of d then $\det(d) = d_1 \cdots d_n$ and $R^n/dR^n = \bigoplus_{j=1}^n R/d_jR$. \square

Note that if $g, g_1 \in M_n(R)$ then $gR^n \subset g_1R^n$ if and only if $g = g_1h$ for some $h \in M_n(R)$, that is if g_1 divides g in $M_n(R)$.

Lemma 2.5. *Let $g \in M_n(R)$. Submodule gR^n is maximal in R^n if and only if $\det(g)$ is nonzero and irreducible in R .*

Proof. Let $g = adb$ be the decomposition from lemma 2.3. Then $\det(g)$ equals to a product of $\det(d)$ and an invertible element of R and R^n/gR^n is isomorphic to R^n/dR^n , thus gR^n is maximal if and only if dR^n is maximal. So we reduced the lemma to the case of diagonal matrices. Let d_1, \dots, d_n be the entries on the diagonal of d . Then $R^n/dR^n = \bigoplus_{j=1}^n R/d_jR$, and therefore dR^n is maximal if and only if one of the d_i is irreducible and the others are invertible, but this happens exactly if $\det d = d_1 \cdots d_n$ is irreducible. \square

Suppose that U is a submodule of R^n . Then by the structure theorem for the finitely generated modules over principal ideal domain, R^n/U is isomorphic to $\bigoplus_{i=1}^n R/q_i^{r_i}R$, where q_i are 0 or irreducible polynomials in $\mathbb{F}_p[x]$ such that $q_i(0) = 1$. Up to permutation, q_i and r_i are uniquely defined by U .

Definition 2.6. Let U be a submodule of R^n , with the quotient R^n/U isomorphic to $\oplus_{i=1}^n R/q_i^{r_i} R$ as above. Define

$$\det^* U = \prod_{i: q_i \neq 0} q_i^{r_i} \in \mathbb{F}_p[x].$$

2.2. Subgroups of \mathcal{A}_n . Let U be an abelian subgroup of \mathcal{A}_n .

Definition 2.7. Define $e(U)$ to be the minimal positive e such that $x^e U = U$, if such e exists, and $+\infty$ otherwise.

Note that if $x^s U = U$ for some $s > 0$ then $e(U)$ divides s .

Lemma 2.8. Define a new structure of left R -module on \mathcal{A}_n by the rule $x * v = x^s v$, denote this new R -module by \mathcal{A}_n^s . Then \mathcal{A}_n^s is isomorphic to R^{ns} .

Proof. Indeed, let e_i , $1 \leq i \leq n$ be the basis of space $(\mathbb{Z}/p\mathbb{Z})^n$ at place 0 in \mathcal{A}_n . Then the map

$$(2.1) \quad R^{ns} \ni (f_{ij})_{1 \leq i \leq n, 0 \leq j \leq s-1} \mapsto \sum_{i,j} f_{ij}(x^s) x^j e_i \in \mathcal{A}_n^s,$$

where $f_{ij} \in R$, is the isomorphism of R modules. \square

Definition 2.9. Suppose that $e(U)$ is finite for some $U \subset \mathcal{A}_n$. By the previous Lemma we can consider U as a submodule of $R^{ne(U)}$. Define $\det^* U$ by the definition 2.6.

We now show that to compute $\det^* U$ it is possible to use other s such that $x^s U = U$.

Proposition 2.10. Suppose $U \subset \mathcal{A}_n$ and $x^s U = U$, and let $f(x) = \det^* U$ where U is considered as a submodule of R^{ns} . Then we also have $x^{sp} U = U$ and if $g(x)$ is $\det^* U$ where U is considered as submodule of R^{nsp} , then $f(x) = g(x)$.

Proof. It follows from the following elementary fact. Suppose $q \in R$ is irreducible, and let M be an R -module $R/q^k R$ with the multiplication $x * (r + q^k R) = x^p r + q^k R$ and $k = tp + b$ for $0 \leq b < p$. Then M is isomorphic to $(R/q^t R)^{p-b} \oplus (R/q^{t+1} R)^b$. \square

2.3. Tree of cosets. To study residually finite groups is the same as to study groups acting faithfully on spherically homogeneous rooted trees (see [14], [17]). There is a bijection between such actions and descending chains of subgroups of finite index with trivial core. The next definition illustrates this.

Definition 2.11. Let G be a group, and $\{H_i | i \geq 1\}$ be the descending chain of subgroups of G , with $H_1 = G$. Construct the following graph: let the vertices be the right cosets of subgroups H_i , that is the set of vertices is $\{gH_i | g \in G, i \geq 1\}$. Let gH_i and $g'H_j$ be connected by an edge iff $i = j + 1$, $gH_i \subset g'H_j$ or $j = i + 1$, $gH_i \supset g'H_j$. Then the constructed graph is a rooted tree with the root $G = H_1$, which is called the tree of cosets corresponding to the chain $\{H_i\}$.

The tree of cosets is locally finite if and only if $(G : H_i) < \infty$ for all i . Note that each $g \in G$ defines an automorphism of the tree of cosets by the multiplication from the right.

The rooted tree is called p -regular if the number of edges going out from each vertex which is not a root is equal to $p + 1$, and the number of vertices going out from the root is equal to p . We denote such a tree by T_p . The group $\text{Aut } T_p$ of all automorphisms of the p -regular tree can be identified with $\wr_1^\infty S_p$, where S_p is the group of all permutations on $\{1, \dots, p\}$. If one considers the action of $\mathbb{Z}/p\mathbb{Z}$ on $\{1, \dots, p\}$ by powers of a standard cycle $(1, \dots, p)$, one can consider $\mathbb{Z}/p\mathbb{Z}$ as a subgroup of S_p . Therefore there is a subgroup of $\text{Aut } T_p$ which is isomorphic to $\wr_1^\infty(\mathbb{Z}/p\mathbb{Z})$. In fact, it is the Sylow p -subgroup of $\text{Aut } T_p$ as a profinite group, see [17]. In the next Proposition we give the following characterization when the tree of cosets is T_p , and when the action of G on it lies in $\wr_1^\infty(\mathbb{Z}/p\mathbb{Z})$.

Proposition 2.12. The tree of cosets is p -regular if and only if the chain is a p -chain, that is $[H_i : H_{i+1}] = p$ for all $i \geq 1$. If the tree of cosets is p -regular then automorphisms of the tree defined by elements of G lie in $\wr_1^\infty(\mathbb{Z}/p\mathbb{Z})$ if and only if the chain is subnormal, that is H_{i+1} is normal in H_i for each $i \geq 1$.

Proof. The first statement is obvious. For the second statement note that the fact that automorphisms of the tree defined by elements of G lie in $\wr_1^\infty(\mathbb{Z}/p\mathbb{Z})$ is equivalent to the fact that if for some $a, g \in G$ and $i \geq 1$ we have that $agH_i = gH_i$ then a acts on cosets of H_{i+1} that lie in gH_i trivially or as some fixed full cycle. Note that since $g^{-1}ag \in H_i$ this is equivalent to the fact that for each $i \geq 1$ the action of H_i on H_i/H_{i+1} by right multiplication acts on H_i/H_{i+1} as powers of some fixed full cycle. Since p is prime this is equivalent to H_{i+1} being normal in H_i . \square

Note that if $\{H_i\}$ is a descending subnormal p -chain then it follows that the stabilizers of the levels of the coset tree are of index a power of p in G .

2.4. Profinite and pro- p topology. For convenience of the reader we remind some well known definitions. Let G be a group.

Definition 2.13. \hat{G} , which is called the profinite closure of G , is the inverse limit of the inverse system $\{G/N\}_N$ where N runs through all normal subgroups of G with finite index, considered with the projective topology, which is called profinite topology. We denote by π the natural homomorphism $\pi : G \rightarrow \hat{G}^{(p)}$.

Note that π is injective if and only if G is residually finite. If in this definition we substitute finite index with index a power of p , we obtain a definition of pro- p closure.

Definition 2.14. $\hat{G}^{(p)}$, called the pro- p closure of G , is the inverse limit of the inverse system $\{G/N\}_N$ where N runs through all normal subgroups of G with index a power of p , considered with the projective topology, which is called pro- p topology. We denote by π_p the natural homomorphism $\pi_p : G \rightarrow \hat{G}^{(p)}$.

Note π_p is injective if and only if G is residually p -finite (i.e. approximated by finite p -groups).

Definition 2.15. Let H be a subgroup of G . We say that H is closed in profinite (respectively pro- p) topology if $\pi(H)$ ($\pi_p(H)$) is equal to the intersection of the closure of $\pi(H)$ with \hat{G} (respectively $\pi_p(H)$ with $\hat{G}^{(p)}$).

We have the following characterization:

Proposition 2.16. Suppose G is finitely generated. A subgroup H of G is closed in pro- p topology if and only if there is a descending subnormal p -chain of subgroups H_i such that $H_1 = G$, $[H_i : H_{i+1}] = p$ for all $i \geq 1$, and $H = \cap_i H_i$.

Proof. Since G is finitely generated we may choose a decreasing sequence N_j of normal subgroups of index a power of p such that $\hat{G}^{(p)} = \varprojlim_j G/N_j$. Then H is closed in pro- p topology iff $H = \cap_j H N_j$, and since $H N_j / N_j \subset H N_{j-1} / N_j$ which is a finite p -group, we can construct a descending subnormal p -chain between $H N_j$ and $H N_{j-1}$ for all j .

Now suppose $H = \cap_i H_i$, $G = H_1$ and $[H_i : H_{i+1}] = p$ and H_{i+1} is normal in H_i for $i \geq 1$. Construct the coset tree which corresponds to it. G acts on it by multiplication from the right, and H is the stabilizer of the ray $\{H_i\}_i$. Let M_i be the stabilizer of i th level of the tree. Then M_i are normal and of index a power of p in G by the proposition. Moreover, $H = \cap_i H M_i$, which implies that $H = \cap_j H N_j$ (since N_j define the pro- p topology, thus for each i there is j such that $M_i \supset N_j$). \square

Remark 1. *Note that a 2-chain is always subnormal. This is the only case when in this article even prime play a special role.*

3. STRUCTURE OF SUBGROUPS

Now we again identify \mathcal{A}_n with the additive group of the ring R^n and convert it into the left R -module with the action $x : v \mapsto xv$. Geometrically one can view this action as given by a left shift, identifying Laurent polynomials with the bilateral sequences over alphabet $\{0, 1, \dots, p-1\}$ (containing only finitely many non-zero entries).

Let us introduce the notation $\varphi_t(x) = 1 + x + \dots + x^{t-1}$. Throughout the article we will use

Lemma 3.1. *Let V be a subgroup of \mathcal{L}_n . Then it defines the triple (s, V_0, v) , where $s \in \mathbb{N}$ is such that $s\mathbb{Z}$ is the image of projection of V on \mathbb{Z} , $V_0 = V \cap \mathcal{A}_n$, satisfying $x^s V_0 = V_0$, and $v \in \mathcal{A}_n$ is such that $(v, s) \in V$. The v is uniquely defined up to addition of elements from V_0 . For $s = 0$ one can choose $v = 0$.*

Conversely any triple (s, V_0, v) with such properties gives rise to a subgroup of \mathcal{L}_n . Two triples (s, V_0, v) and (s', V'_0, v') define the same subgroup if and only if $s = s'$, $V_0 = V'_0$ and $v + V_0 = v' + V_0$.

Moreover, $V \subset V'$ if and only if $s'|s$, $V_0 \subset V'_0$ and $v = \varphi_{s/s'}(x^{s'})v' \bmod V'_0$.

Proof. To show the first part of the Lemma we need only to prove that $x^s V_0 = V_0$. Indeed,

$$(v, s)(w, 0)(-x^{-s}v, -s) = (x^s w, 0) \in V,$$

and so $x^s w \in V \cap \mathcal{A}_n = V_0$. Thus $x^s V_0 \subset V_0$. The converse inclusion is proved in a similar way.

Now suppose we are given a triple (s, V_0, v) such that V_0 is a subgroup of \mathcal{A}_n and $x^s V_0 = V_0$. Let V be the subgroup of \mathcal{L}_n generated by V_0 and (v, s) . To show that the triple (s, V_0, v) is a triple for V we need to show that the projection of V onto \mathbb{Z} is $s\mathbb{Z}$ and that $V \cap \mathcal{A}_n = V_0$. Since $V_0 \subset \mathcal{A}_n$ the projection of V on \mathbb{Z} is indeed $s\mathbb{Z}$. Clearly, $V_0 \subset V \cap \mathcal{A}_n$. To prove the converse inclusion note first that if $s = 0$ then $v \in V_0$ and therefore $V = (V_0, 0)$. Suppose that $s \geq 1$. Then by the equality above any element of V is of the form $(w, 0)(v, s)^k$ for some $w \in V_0$ and $k \in \mathbb{Z}$, and $(w, 0)(v, s)^k \in V_0$ if and only if $k = 0$, which implies that $V_0 = V \cap \mathcal{A}_n$.

Suppose now that we have two triples (s, V_0, v) and (s', V'_0, v') , and V is the subgroup generated by V_0 and (v, s) , while V' is the subgroup generated by V'_0 and (v', s') . It is left to prove that $s = s'$, $V_0 = V'_0$, and $v + V_0 = v' + V_0$. First two equalities immediately follow from the

definitions of s and V_0 in terms of V . On the other hand we know that (v, s) and (v', s) both belong to $V = V'$. It follows that

$$(v', s)(v, s)^{-1} = (v', s)(-x^{-s}v, -s) = (v' - v, 0)$$

also belongs to V . Thus $v' - v \in V_0$ and therefore $v + V_0 = v' + V_0$.

To prove the last part, note that both (v, s) and $(\varphi_{s/s'}(x^{s'})v', s) = (v', s')^{s/s'}$ belong to V' , and therefore $v = \varphi_{s/s'}(x^{s'})v' \bmod V'_0$. Conversely, if $s|s'$, $V_0 \subset V'_0$ and $v = \varphi_{s/s'}(x^{s'})v' + w$ for some $w \in V'_0$, then $(v, s) = (w, 0)(v', s')^{s/s'} \in V'$, and therefore $V \subset V'$. \square

Remark 2. Subgroup V belongs to \mathcal{A}_n if and only if $s = 0$ in the triple corresponding to V .

We have the description of normal subgroups of \mathcal{L}_n in terms of the triple.

Lemma 3.2. Let V be a subgroup of \mathcal{L}_n . Then V is normal if and only if the corresponding triple (s, V_0, v) , defined in Lemma 3.1 satisfies the additional properties that $xV_0 = V_0$, $(1 - x^s)\mathcal{A}_n \subset V_0$, and $(1 - x)v \in V_0$.

Proof. Take $w \in V_0$. Then, both

$$(0, 1)(w, 0)(0, -1) = (xw, 0),$$

$$(0, -1)(w, 0)(0, 1) = (x^{-1}w, 0),$$

belong to V since V is normal. Thus both xw and $x^{-1}w$ also belong to V_0 . It follows that $xV_0 = V_0$.

Take now $w \in \mathcal{A}_n$. Then $(w, 0)(v, s)(-w, 0) = ((1 - x^s)w + v, s) \in V$, since $(v, s) \in V$ and V is normal. It follows that $(1 - x^s)w + v + V_0 = v + V_0$, thus $(1 - x^s)w \in V_0$, and it is true for any $w \in \mathcal{A}_n$. In a similar way $(0, 1)(v, s)(0, -1) = (xv, s) \in V$, therefore $v + V_0 = xv + V_0$, and thus $(1 - x)v \in V_0$. The proof of the converse statement, that if (s, V_0, v) satisfies such additional properties then V is normal, is also straightforward. \square

Corollary 3.3. Let V is a normal subgroup of \mathcal{L}_n with the triple (s, V_0, v) . Then if $s > 0$ it has finite index in \mathcal{L}_n . If $n = 1$ and V is nontrivial, V_0 has finite index in \mathcal{A}_n .

Proof. If $s > 0$ then $(x^s - 1)\mathcal{A}_n$ has finite index in \mathcal{A}_n , therefore since $(x^s - 1)\mathcal{A}_n \subset V_0$, we obtain that V_0 also has finite index in \mathcal{A}_n and hence V has finite index in \mathcal{L}_n .

Let $n = 1$. Then if $s > 0$ we are done, so suppose $s = 0$ and $V = V_0 \subset \mathcal{A}_1$. By the previous lemma $xV_0 = V_0$, hence identifying \mathcal{A}_1 with R we obtain that V_0 is the nontrivial ideal of R . Any such ideal is of the form fR for some $f \in \mathbb{F}_p[x]$ and hence is of finite index in R . \square

Corollary 3.4. *Let V, W be the subgroups of \mathcal{L}_n , with corresponding triples (s, V_0, v) and (t, W_0, w) . Let $U = V \cap W$. Choose $r > 0$ minimal with the property that both s and t divide r and moreover*

$$\frac{1-x^r}{1-x^s}v = \frac{1-x^r}{1-x^t}w \pmod{(V_0 + W_0)},$$

and if no such number exists put $r = 0$.

If $r = 0$ then $V \cap W = V_0 \cap W_0 \subset \mathcal{A}_n$, in particular $V \cap W = \{0\}$ if $r = 0$ and $V_0 \cap W_0 = \{0\}$.

If $r > 0$ then exist $v_0 \in V_0$ and $w_0 \in W_0$ such that

$$\frac{1-x^r}{1-x^s}v + v_0 = \frac{1-x^r}{1-x^t}w + w_0.$$

Denote this element by u . Then $(r, V_0 \cap W_0, u)$ is the triple that corresponds to the subgroup U .

Proof. Observe that $u = \varphi_{r/s}(x^s)v + v_0$ and that $(1-x^r)/(1-x^s) = \varphi_{r/s}(x^s)$. Assume $r = 0$. Suppose there is $g = (x, q) \in V \cap W$ with $q \neq 0$. It follows that $(x, q) = (v_0, 0)(v, s)^n$ for some $v_0 \in V_0$ and integer n . But $(v, s)^n = (\varphi_n(x^s)v, sn)$ and therefore $x = v_0 + \varphi_n(x^s)v$ and $q = sn$. Analogously since $g = (x, q) \in W$ it follows that $q = tm$ and $x = w_0 + \varphi_m(x^t)w$ for some $w_0 \in W_0$ and some integer m . Therefore $\varphi_{q/s}(x^s)v = \varphi_{q/t}(x^t)w \pmod{(V_0 + W_0)}$ and thus $r > 0$, contradiction. It follows that for any $g \in V \cap W$ the projection onto \mathbb{Z} is 0, thus $V \cap W \subset \mathcal{A}_n$.

Let now $r > 0$. The subgroup defined by the triple $(r, V_0 \cap W_0, u)$ belongs both to V and W . Indeed, we just need to prove that $u = \varphi_{r/s}(x^s)v \pmod{V_0}$ and $u = \varphi_{r/t}(x^t)w \pmod{W_0}$, but this directly follows from the definition of r and u .

Conversely, suppose that a subgroup U' with the triple (r', U'_0, u') belongs to both V and W . We need to prove that $U' \subset U$, that is, by the Lemma, that $U'_0 \subset V_0 \cap W_0$, $r|r'$ and $u' = \varphi_{r'/r}(x^r)u \pmod{V_0 \cap W_0}$.

Since $u' = \varphi_{r'/s}(x^s)v \pmod{V_0}$ and $u' = \varphi_{r'/t}(x^t)w \pmod{W_0}$ it follows that $\varphi_{r'/s}(x^s)v = \varphi_{r'/t}(x^t)w \pmod{V_0 + W_0}$, therefore by the minimality of r $r \leq r'$. Let $r' = ar + b$ with $b < r$. Then

$$\begin{aligned} \varphi_{r'/s}(x^s)v &= \frac{1-x^{r'}}{1-x^s}v = \frac{1-x^b}{1-x^s}v + x^b \frac{1-x^{ar}}{1-x^r} \varphi_{r/s}(x^s)v, \\ \varphi_{r'/t}(x^t)w &= \frac{1-x^{r'}}{1-x^t}w = \frac{1-x^b}{1-x^t}w + x^b \frac{1-x^{ar}}{1-x^r} \varphi_{r/t}(x^t)w, \end{aligned}$$

and thus

$$\frac{1-x^b}{1-x^s}v = \frac{1-x^b}{1-x^t}w \pmod{V_0 + W_0}.$$

Since $b < r$ we obtain that $b = 0$ and therefore r divides r' .

To show that $u' = \varphi_{r'/r}(x^r)u \pmod{V_0 \cap W_0}$ note that

$$\varphi_{r'/r}(x^r)u = \frac{1 - x^{r'}}{1 - x^r}u = \frac{1 - x^{r'}}{1 - x^s}v + \varphi_{r'/r}(x^r)v_0 = \frac{1 - x^{r'}}{1 - x^t}w + \varphi_{r'/r}(x^r)w_0,$$

and since r is divisible by both s and t , $\varphi_{r'/r}(x^r)v_0 \in V_0$ and $\varphi_{r'/r}(x^r)w_0 \in W_0$. Therefore $u' = \varphi_{r'/r}(x^r)u \pmod{V_0}$ and $u' = \varphi_{r'/r}(x^r)u \pmod{W_0}$, which finishes the proof. \square

The structure of subgroups of \mathcal{L}_n can be described as follows.

Theorem 3.5. *Suppose V is a subgroup of \mathcal{L}_n . Then either V is a subgroup of \mathcal{A}_n or V is isomorphic to \mathcal{L}_k for some $k \in \mathbb{N}$. In particular if V is of finite index in \mathcal{L}_n then V is isomorphic to \mathcal{L}_{ns} where s is the projection of V onto \mathbb{Z} .*

Proof. Let (s, V_0, v) be some triple corresponding to V . If $s = 0$ then $V = V_0$ is a subgroup of \mathcal{A}_n . Suppose that $s \geq 1$. Recall (Lemma 2.8) that we can consider \mathcal{A}_n as a left R -module, which we denoted \mathcal{A}_n^s , by defining the action of x on v as $x * v = x^s v$, $v \in \mathcal{A}_n$. Then V_0 is R submodule. By Lemma 2.8 \mathcal{A}_n^s is a free R module of rank ns . Since R is a principal ideal domain, it is noetherian, and thus V_0 is finitely generated as R module, and therefore, as a submodule of the free finitely generated R module R^{ns} V_0 is also free, i.e. V_0 is isomorphic to R^k for some k . V_0 is of finite index in \mathcal{A}_n iff $k = ns$. Now let H be the subgroup of V generated by (v, s) . Then V_0 is normal in V , $V_0 \cap H = \{0\}$, H is isomorphic to \mathbb{Z} . It follows that V is isomorphic to $V_0 \rtimes \mathbb{Z}$, where the action of the generator of \mathbb{Z} on V is given by the map x^s . It follows, by above, that V is isomorphic to $R^k \rtimes \mathbb{Z}$ where \mathbb{Z} acts on R^k by multiplication by x . Therefore, V is isomorphic to \mathcal{L}_k . If V is of finite index in \mathcal{L}_n then $k = ns$. \square

Corollary 3.6. *Any finitely generated subgroup of \mathcal{L}_n is closed in the profinite topology.*

This result also follows from Proposition 3.19 in [10]. Groups with stated property are called LERF or subgroup separable.

Proof. Let V be a finitely generated subgroup of \mathcal{L}_n and let (s, V_0, v) be the corresponding triple. If $s = 0$ then $V \subset \mathcal{A}_n$, and since it is finitely generated, it is finite, hence closed. If $s > 0$, then V belongs to the subgroup W with the triple $(s, \mathcal{A}_n, 0)$. W has finite index in \mathcal{L}_n , and therefore V is profinitely closed in \mathcal{L}_n if and only if it is closed in W . Moreover, W is isomorphic to \mathcal{L}_{ns} , and V , considered as a subgroup of \mathcal{L}_{ns} , has the projection on \mathbb{Z} equal to \mathbb{Z} . Hence we have reduced to the case when $s = 1$. For any $v \in \mathcal{A}_n$, there is an automorphism φ of \mathcal{L}_n which is identical on \mathcal{A}_n and $\varphi(0, 1) = (v, 1)$. Since the property of

being profinitely closed is preserved under an automorphism, and the triple for $\varphi(V)$ is $(1, V_0, 0)$, we have reduced to the case when $s = 1$ and $v = 0$. Changing the basis in \mathcal{A}_n we reduce to the case when $V_0 = \bigoplus_{i=1}^n f_i R$, which is closed in the profinite topology. \square

Corollary 3.7. *Any subgroup of index p of \mathcal{L}_n is isomorphic either to \mathcal{L}_n or to \mathcal{L}_{pn} .*

Proof. Indeed, if V is of index p then the corresponding s defined in Lemma 3.1 is either 1 or p . By the previous Lemma, if $s = 1$ then V is isomorphic to \mathcal{L}_n , and if $s = p$ then V is isomorphic to \mathcal{L}_{pn} . \square

3.1. Subgroup Growth. Now we compute the number of subgroups of given index of \mathcal{L}_n . First let us compute the number of submodules of R^k of given finite codimension. Define

$$b_t(R^k) = |\{M \subset R^k : M \text{ is } R \text{ submodule and } \dim_{\mathbb{F}_p} R^k/M = t\}|.$$

Lemma 3.8. $b_t(R^k) = p^{tk} - p^{(t-1)k}$ for $t > 0$, $b_0(R^k) = 1$.

Proof. We use a similar method that was used in the case of \mathbb{Z}^k for counting subgroup growth (see for instance [27]). It is based on consideration of endomorphisms of R^k . Since R is principal ideal domain, every submodule of R^k is isomorphic to R^s , $s \leq k$, and if it is of finite index, then $s = k$. Therefore, it is equal to gR^k , for some $g \in M_k(R)$, $\det(g) \neq 0$. Moreover, the index of gR^k in R^k is equal to the number of elements in $R/(\det g)$. Also, $g_1R^k = g_2R^k$ if and only if $g_2^{-1}g_1 \in GL_k(R)$. It is easy to see that the orbit representatives of the action by $GL_k(R)$ by the multiplication from the right on the set $\{g \in M_k(R) | \det(g) \neq 0\}$ are lower triangular matrices (g_{ij}) such that $g_{ij} \in \mathbb{F}_p[x]$ for all i, j , $g_{ii}(0) = 1$ for all i , $g_{ij} = 0$ for $i < j$, and the degree of g_{ij} is less than g_{jj} for all i, j .

Introduce function ϕ by $\phi(s) = (p-1)p^{s-1}$ if $s \geq 1$ and $\phi(0) = 1$. Note that $\sum_{s=1}^n \phi(s) = p^n$ and that the number of polynomials $g \in \mathbb{F}_p[x]$ such that $g(0) = 1$ and the degree of g is s is equal to $\phi(s)$. Then the number of all submodules of codimension t (that is, of index p^t) of R^k is equal to the number of the lower triangular matrices described above with sum of degrees $\sum_i \deg(g_{ii}) = t$, that is to

$$\sum_{x_1 + \dots + x_k = t} \phi(x_1) \dots \phi(x_k) p^{x_2 + \dots + (k-1)x_k}.$$

To compute it we first compute

$$\begin{aligned}
 & \sum_{x_1+\dots+x_k \leq t} \phi(x_1) \cdots \phi(x_k) p^{x_2+\dots+(k-1)x_k} = \\
 (3.1) \quad & \sum_{x_2+\dots+x_k \leq t} \phi(x_2) \cdots \phi(x_k) p^{x_2+\dots+(k-1)x_k} \sum_{x_1=0}^{t-(x_2+\dots+x_k)} \phi(x_1) = \\
 & p^t \sum_{x_2+\dots+x_k \leq t} \phi(x_2) \cdots \phi(x_k) p^{x_3+\dots+(k-2)x_k} = \dots = p^{kt},
 \end{aligned}$$

therefore $b_t(R^k) = p^{tk} - p^{(t-1)k}$

□

Proposition 3.9. *Let $a_m(\mathcal{L}_n)$ be the number of subgroups of \mathcal{L}_n of index m . Then*

$$a_m(\mathcal{L}_n) = \sum_{t: p^t | m} b_t(R^{\frac{nm}{p^t}}) p^t = \sum_{t \geq 1: p^t | m} (p^{\frac{nm}{p^t}} - p^{\frac{nm(t-1)}{p^t}}) p^t + 1.$$

Proof. If V is the subgroup of \mathcal{L}_n and (s, V_0, v) is the corresponding triple, then its index is equal to $s[\mathcal{A}_n : V_0]$. The number of subgroups with fixed s and V_0 is equal to number of coset representatives v , i.e. to the index $[\mathcal{A}_n : V_0]$. The formula now follows. □

Proposition 3.10. *Let $s_m(\mathcal{L}_n) = \sum_{1 \leq d \leq m} a_d(\mathcal{L}_n)$. Then*

$$s_m(\mathcal{L}_n) \geq p(p^{n[m/p]+1} - 1),$$

where $[m/p]$ is the integer part of m/p .

Proof. First we write $s_m(\mathcal{L}_n)$ as

$$\begin{aligned}
 s_m(\mathcal{L}_n) = & m + \sum_{t, k \geq 1: p^t k \leq m} (p^{nkt} - p^{nk(t-1)}) p^t = m + \sum_{t=1}^{[\log_p m]} p^t \sum_{k=1}^{[m/p^t]} (p^{nkt} - p^{nk(t-1)}) = \\
 & m - p \left[\frac{m}{p} \right] + \frac{p^{n+1}}{p^n - 1} [p^{n[m/p]} - 1] - \frac{p^{n+2}}{p^n - 1} [p^{n[m/p^2]} - 1] + \\
 & \frac{p^{2n+2}}{p^{2n} - 1} [p^{2n[m/p^2]} - 1] + \sum_{t=3}^{[\log_p m]} p^t \sum_{k=1}^{[m/p^t]} (p^{nkt} - p^{nk(t-1)}),
 \end{aligned}$$

where

$$\sum_{t=3}^{[\log_p m]} p^t \sum_{k=1}^{[m/p^t]} (p^{nkt} - p^{nk(t-1)}) \leq \frac{p^{3n}}{p^{3n} - 1} p^{3(nm/p^3+1)} \log_p m$$

□

Remark 3. Note that for $n = 1$ this bound gives a slight improvement over the bound in proposition 3.2.1 in [27], where it was shown that $s_m(\mathcal{L}_1) > c^m$ for $c < p^{1/p^2}$ and big enough m .

Now we are going to consider the structure of maximal and weakly maximal subgroups of \mathcal{L}_n (where by weakly maximal we understand those subgroups of infinite index such that any larger subgroup is of finite index).

Theorem 3.11. *Each maximal subgroup has finite index and is either of the form $\mathcal{A}_n \rtimes q\mathbb{Z}$ for some prime $q \in \mathbb{Z}$, in which case its index is q , or its corresponding triple is $(1, V_0, v)$ such that $\det^* V_0$ is irreducible polynomial, in which case its index is $p^{\deg(\det^* V_0)}$. Maximal subgroups of the form $\mathcal{A}_n \rtimes q\mathbb{Z}$ are normal. Maximal subgroup with the triple $(1, V_0, v)$ is normal if and only if $\det^* V_0 = 1 - x$.*

Proof. Suppose V is a maximal subgroup and (s, V_0, v) is the corresponding triple. Then there are three possible cases. If $s = 0$ then $V = V_0 \subset \mathcal{A}_n$, and since it is maximal, it must be equal to \mathcal{A}_n . However, \mathcal{A}_n is not maximal, as it is contained in, say, $\mathcal{A}_n \rtimes 2\mathbb{Z}$.

If $s > 1$ then, by maximality $V_0 = \mathcal{A}_n$, thus v can be taken to be 0, therefore the group is $\mathcal{A}_n \rtimes s\mathbb{Z}$. If q is any prime divisor of s then $\mathcal{A}_n \rtimes s\mathbb{Z}$ is contained in $\mathcal{A}_n \rtimes q\mathbb{Z}$, therefore for maximal subgroups s is prime. On the other hand if a subgroup with triple $(q, \mathcal{A}_n, 0)$ is contained in the nontrivial subgroup with triple (t, W, w) then clearly $W = \mathcal{A}_n$ and s divides q . Since q is prime, $s = q$ or $s = 1$. By nontriviality $s = q$ is the only choice. Thus $\mathcal{A}_n \rtimes q\mathbb{Z}$ is maximal for any prime q . It is clear that its index is equal to $(\mathbb{Z} : q\mathbb{Z}) = q$.

Finally let $s = 1$ and $(1, V_0, v)$ be the corresponding triple for a maximal subgroup V . By the lemma above $xV_0 = V_0$. By considering \mathcal{A}_n as a module over R isomorphic to R^n we obtain that V_0 is its submodule, and therefore isomorphic to gR^n for some $g \in M_n(R)$. Moreover V_0 is maximal submodule, by the maximality of the subgroup V . By Lemma 2.5, it happens if and only if $\det g$ is irreducible. Notice then that $\det^* V_0 = u \det(g)$ for some invertible element $u \in R$. In this case the index of V in \mathcal{L}_n is equal to the index of V_0 in \mathcal{A}_n , which is equal to $p^{\deg(\det g)} = p^{\deg(\det^* V_0)}$ by the corollary after Lemma 2.3.

It is obvious that $\mathcal{A}_n \rtimes q\mathbb{Z}$ are normal. To consider when a subgroup with the triple $(1, V_0, v)$ such that V_0 is isomorphic to gR^n is normal, let $g = adb$ be the product from Lemma 2.3. Then $\det^* V_0 = d_1 \cdots d_n$, and it is irreducible iff for some i d_i is irreducible and all other $d_j = 1$. The group with a triple $(1, gR^n, v)$ is normal iff $(1 - x)R^n \subset gR^n$, since $(1 - x)v \in gR^n$ follows from it. But since $gR^n = adbR^n = adR^n$, the condition $(1 - x)R^n \subset gR^n$ is equivalent to the condition $(1 - x)R^n \subset$

dR^n , that is $(1-x)R \subset d_k R$ for all k , hence each d_k must divide $x-1$. Therefore the maximal subgroup with the triple $(1, gR^n, v)$ is normal iff $\det(g) = u(x-1)$ for some invertible element u of R . \square

Theorem 3.12. *Let V be a subgroup of \mathcal{L}_n , and (s, V_0, v) be its triple. Then V is weakly maximal if and only if either $s = 0$ and $V_0 = \mathcal{A}_n$, or $s > 0$ and the factormodule \mathcal{A}_n/V_0 is isomorphic to R as R modules under the identification of \mathcal{A}_n and R^{ns} from Lemma 2.8. Such weakly maximal subgroup V is closed in pro- p topology if and only if $s = 0$ or s is a power of p .*

Proof. First let us prove that \mathcal{A}_n is weakly maximal. Indeed, it is of infinite index and any larger subgroup must have nontrivial projection onto \mathbb{Z} , i.e. its corresponding triple is $(s, \mathcal{A}_n, 0)$, with $s \geq 1$ and hence has finite index s .

Suppose V is a weakly maximal subgroup and let (s, V_0, v) be the corresponding triple. If $s = 0$ then $V = V_0 \subset \mathcal{A}_n$, and since V is weakly maximal we have equality $V = \mathcal{A}_n$. If $s \geq 1$ then \mathcal{A}_n^s/V_0 is isomorphic as R module to $R/(g_1) \oplus \cdots \oplus R/(g_k) \oplus R^t$, and by weak maximality of V it follows that $k = 0$ and $t = 1$. Conversely, suppose that V is such a subgroup that \mathcal{A}_n^s/V_0 is isomorphic to R as modules over R . Suppose it is not weakly maximal, i.e. there is a subgroup V' of infinite index that contains V . Let (s', V'_0, v') be the corresponding triple for V' . Then s' divides s and $V_0 \subset V'_0$. Since s is a multiple of s' we have that $x^{s'}V'_0 = V_0$, thus V'_0 is an R submodule of \mathcal{A}_n^s that contains V_0 . Then either V'_0 is equal to V_0 or $\mathcal{A}_n^s/V'_0 = R/(g')$ for some nonzero g' . But in the latter case V'_0 is of finite index in \mathcal{A}_n and therefore V' is of finite index in \mathcal{L}_n , which contradicts our assumption. Thus $V'_0 = V_0$. It is left to prove that $s = s'$. Let $s = s'k$. Let M be the factor \mathcal{A}_n/V_0 . Since $x^{s'}V_0 = V_0$, we may consider M as a module over R , with multiplication by x given by the action of $x^{s'}$. To show that $k = 1$ and therefore $s = s'$ it is left to prove the following lemma:

Lemma 3.13. *Let M be the module over R , and let $\pi : R \rightarrow R$ be the ring homomorphism given by $\pi(x) = x^k$. Consider new R module M_π with the same underlying space as M and with the action of R given by $r * m = \pi(r)m$. Suppose that M_π is isomorphic to R as modules over R . Then $k = 1$.*

Proof. Since M_π is isomorphic to R as modules over R there is $m \in M$ such that the map, $r \mapsto \pi(r)m$ is bijective. Consider now map $\eta : R \rightarrow M$, $r \mapsto rm$, and let $R' = \pi(R) \subset R$. Then η is linear over \mathbb{F}_p and its restriction to R' is bijective. It follows that $R = R'$ and thus $k = 1$. \square

Note that we have actually shown that if V is weakly maximal and the corresponding s is nonzero, then it is the smallest positive integer such that V_0 is invariant under multiplication by x^s .

Now let V be a weakly maximal subgroup which is closed in pro- p topology, and suppose $s > 0$. By the Proposition 2.16 there is a descending subnormal p -chain $\{H_i\}$ such that $V = \cap_i H_i$, and let (s_i, V_i, v_i) be a triple associated to H_i . Then $s_i | s_{i+1}|s$ for all i , hence the sequence s_i must stabilize. Thus $s_i = p^t$ for some t and all big enough i . Since $V = \cap_i H_i$, we obtain $V_0 = V \cap \mathcal{A}_n = \cap_i V_i$. The fact that $x^{p^t} V_i = V_i$ for all i implies that $x^{p^t} V_0 = V_0$, and since s is smallest such integer we must have $s = p^t$.

Conversely, suppose that V is weakly maximal and $s = p^t$, then $V \subset \mathcal{A}_n \rtimes p^t \mathbb{Z}$. There is a subnormal p -chain $\mathcal{A}_n \rtimes p^j \mathbb{Z}$, $0 \leq j \leq t$ between $\mathcal{A}_n \rtimes p^t \mathbb{Z}$ and \mathcal{L}_n , thus it is left to construct the subnormal p -chain between V and $\mathcal{A}_n \rtimes p^t \mathbb{Z}$. Identifying $\mathcal{A}_n \rtimes p^t \mathbb{Z}$ with \mathcal{L}_{np^t} we reduce to the case $t = 0$. Now, since V is weakly maximal, \mathcal{A}_n/V_0 is isomorphic to R as R modules, and denote the corresponding map $\mathcal{A}_n \rightarrow R$ by ψ . Then $V_i = \psi^{-1}((1+x)^i R)$ form a p -chain between V_0 and \mathcal{A}_n , and hence groups with triples $(1, V_i, v)$ form subnormal p -chain between V and \mathcal{L}_n , where v is a vector from a triple of V . Hence there is a p -chain that intersects to V and thus V is closed in pro- p topology by the Proposition 2.16. \square

We recall the following definition.

Definition 3.14. (see [15]) *The group is called hereditary free from finite normal subgroups if any subgroups of it of finite index have only infinite nontrivial normal subgroups.*

Corollary 3.15. *Any nontrivial normal subgroup of \mathcal{L}_n is infinite. Therefore, \mathcal{L}_n is hereditary free from finite normal subgroups.*

Therefore we can derive from [15] the following result:

Corollary 3.16. *Any faithful action of \mathcal{L}_n on the rooted tree induces topologically free action on the boundary of the tree.*

3.2. Growth of normal subgroups of \mathcal{L}_n . Recall that the zeta function for normal subgroups of \mathcal{L}_n is the following:

$$(3.2) \quad \zeta_n(z) = \sum_{N \triangleleft \mathcal{L}_n : [\mathcal{L}_n : N] < \infty} [\mathcal{L}_n : N]^{-z}$$

We have the following formula:

Theorem 3.17.

$$\zeta_n(z) = \sum_{s \geq 1} s^{-z} \prod_{r \in R, \text{prime}, r|1-x^s, r \neq 1-x} \left(\sum_{M: M \in \text{Mod}_r, (1-x^s)R^n \subset M} [R^n : M]^{-z} \right) \times$$

$$\left(\sum_{M: M \in \text{Mod}_{1-x}, (1-x^s)R^n \subset M} \tau(M) [R^n : M]^{-z} \right),$$

where Mod_r is the set of all submodules M in R^n such that $r^k R^n \subset M$ for some k , and, for $M \in \text{Mod}_{1-x}$, $\tau(M) = |\ker \varphi|$ for $\varphi : R^n/M \rightarrow R^n/M$, $\varphi(r) = (1-x)r$.

Proof. Recall that if M is a submodule of R^n of finite index, then $R^n/M = \bigoplus_{r \in R, r \text{ prime}} N_r$ is the primary decomposition of R^n/M (Theorem 6.7 in [22]), with only finite number of N_r not zero. Let $\pi : R^n \rightarrow R^n/M$ and let $M_r = \pi^{-1}(N_r)$. Then $M_r \in \text{Mod}_r$, $M = \bigcap_r M_r$, and $[R^n : M] = \prod_r [R^n : M_r]$. We obtain

$$\zeta_n(z) = \sum_{s \geq 1, M \subset R^n, v \in R^n/M : (1-x^s)R^n \subset M, (1-x)v=0} (s[R^n : M])^{-z} =$$

$$\sum_{s \geq 1} s^{-z} \prod_{r \in R, \text{prime}, r|1+x^s, r \neq 1-x} \left(\sum_{M: M \in \text{Mod}_r, (1-x^s)R^n \subset M} [R^n : M]^{-z} \right) \times$$

$$\left(\sum_{M: M \in \text{Mod}_{1-x}, (1-x^s)R^n \subset M} \tau(M) [R^n : M]^{-z} \right)$$

□

Corollary 3.18.

$$\zeta_n(z) \preceq p^n \zeta(z) \xi_n(z)$$

where

$$\zeta(z) = \sum_{s \geq 1} s^{-z}$$

is the Riemann zeta function,

$$\xi_n(z) = \sum_{M \subset_f R^n} [R^n : M]^{-z}$$

where the summation is over submodules of finite index in R^n , is the zeta function of the module R^n , and

$$\sum_{m \geq 1} b_m m^{-z} \preceq \sum_{m \geq 1} c_m m^{-z}$$

if and only if $0 \leq b_m \leq c_m$ for all $m \geq 1$.

Proof. Note that for $M \in \text{Mod}_{1-x}$, we have that $R^n/M = \oplus_1^n R/(1-x)^{k_i} R$, therefore $\dim_{\mathbb{F}_p} \ker \phi \leq n$ and thus $\tau(M) \leq p^n$. Note also that

$$\xi_n(z) = \sum_{M \subset_f R^n} [R^n : M]^{-z} = \prod_{r \in R, r \text{ prime}} \left(\sum_{M \in \text{Mod}_r} [R^n : M]^{-z} \right),$$

and thus

$$\sum_{M: M \in \text{Mod}_r, (1-x^s)R^n \subset M} [R^n : M]^{-z} \leq \sum_{M \in \text{Mod}_r} [R^n : M]^{-z},$$

and we obtain the first formula. \square

Corollary 3.19. *Let $s_m^\triangleleft(\mathcal{L}_n)$ be the number of normal subgroups of \mathcal{L}_n of index smaller or equal to m . Then*

$$\limsup_m \frac{\log s_m^\triangleleft(\mathcal{L}_n)}{\log m} \leq n$$

Proof. Notice that by the lemma 3.8

$$\xi_n(z) = 1 + \sum_{t \geq 1} \frac{p^{tn} - p^{(t-1)n}}{p^{tz}} = \frac{p^z - 1}{p^z - p^n},$$

so $\xi_n(z)$ is analytic for $\Re z > n$, where $\Re z$ is the real part of z , and thus by the corollary 3.18 $\zeta_n(z)$ is analytic for $\Re z > n$, since Riemann zeta function is analytic for $\Re z > 1$. Now observe that

$$\zeta_n(z) = \sum_{m=1}^{\infty} \frac{a_m^\triangleleft(\mathcal{L}_n)}{m^z},$$

where $a_m^\triangleleft(\mathcal{L}_n)$ is the number of normal subgroups of \mathcal{L}_n of index m , and note that $s_m^\triangleleft(\mathcal{L}_n) = \sum_{j=1}^m a_j^\triangleleft(\mathcal{L}_n)$, from which we obtain the statement of the corollary (see the beginning of chapter 15 of [27]). \square

Remark 4. *Note that to derive the estimate Corollary 3.19 we only used the property that if V is a normal subgroup of \mathcal{L}_n then $V \cap \mathcal{A}$ is invariant under the shift, no other properties of normal subgroups listed in the Lemma 3.2 were used.*

3.3. Subgroups, normal subgroups and factors of $\mathcal{L}_{1,p}$. When $n = 1$ we give one more description of normal subgroups.

Lemma 3.20. *Let $e \in \mathcal{A}_1$ be the element with 1 on the position 0 and zeros everywhere else. Then to each nontrivial normal subgroup V of \mathcal{L}_1 uniquely corresponds triple (s, f, h) where $s \in \mathbb{N}$ is defined as before, and $f, h \in \mathbb{F}_p[x]$ are defined in the following way: f is the polynomial of minimal degree such that $f(x)e \in V$ and $f(0) = 1$, and*

h is the polynomial such that $\deg h < \deg f$ and $h(x)e = v \pmod{V_0}$. Moreover, $f(0) = 1$. If $s \geq 1$ then f divides $1 - x^s$, and h is either 0 or $h = f(1 - x)^{-1}$ (the latter is possible only if f is divisible by $1 - x$).

Proof. Since V is normal and not trivial then V_0 is also not trivial. Indeed if $(v, s) \in V$ then the commutator $[(w, 0), (v, s)] = (w - x^s w, 0) \in V_0$. Note that if we identify \mathcal{A}_1 and R , with e identified to $1 \in R$, then each nontrivial subgroup V_0 of \mathcal{A}_1 such that $xV_0 = V_0$ is in fact a nontrivial ideal of R , which must be of the form fR for some polynomial $f \in \mathbb{F}_p[x]$ such that $f(0) = 1$. Under this identification \mathcal{A}_1/V_0 is identified with $R/fR = \mathbb{F}_p[x]/f\mathbb{F}_p[x]$, and since v from the triple is defined up to addition of element of V_0 , we choose the unique polynomial of the minimal degree h which is the representative of the corresponding coset of $\mathbb{F}_p[x]/f\mathbb{F}_p[x]$.

Now let V be the normal subgroup of \mathcal{L}_1 , (s, V_0, v) , (s, f, h) be the triples associated to it as defined in the statement of the lemma. We already proved that $(1 - x^s)\mathcal{A}_1 \subset V_0$, which implies that $(1 - x^s)e \in V_0$ and thus by the definition of f that $1 - x^s$ is divisible by f . Now, suppose $h \neq 0$. On the one hand the degree of h is less than the degree of f , since otherwise we reduce it preserving the requirements on h . On the other hand it follows from $(1 - x)v \in V_0$ and $v = h(x)e \pmod{V_0}$ that $(1 - x)h(x)e \in V_0$, and thus that f divides $(1 - x)h$. Therefore $f = (1 - x)h$. \square

To summarize, we have the following three cases for the triple (s, f, h) associated to the normal subgroup V of \mathcal{L} : either $s = 0$ and then the triple is $(0, f, 0)$, with no restrictions on f except $f(0) = 1$; or $s \geq 1$, then necessarily $f \neq 0$, and either $h = 0$ and the triple is $(s, f, 0)$ with $f|(1 - x^s)$ or $h \neq 0$ and the triple is $(s, (1 - x)h, h)$ with h dividing $(1 - x^s)(1 - x)^{-1} = 1 + x + \dots + x^{s-1} = \varphi_s(x)$.

Definition 3.21. For $f \in \mathbb{F}_p[x]$ and $s \geq 1$, such that $f(0) = 1$ and $f|1 - x^s$ denote by $B_{f,s}$ the corresponding subgroup with projection onto \mathbb{Z} equal to $s\mathbb{Z}$, intersection with base equal to fR and the corresponding vector equal to 0. If moreover $1 - x|f$, that is if $f(1) = 0$, denote the group with triple $(s, f, f(1 - x)^{-1})$ by $C_{f,s}$.

Theorem 3.22. Any nontrivial normal subgroup of \mathcal{L}_1 of finite index is either $B_{f,s}$ or $C_{f,s}$ for some polynomial f and $s \geq 1$. We have the following description of the lattice of normal subgroups of finite index in \mathcal{L}_1 .

- $B_{f,s} \subset B_{f',s'}$ if and only if $s'|s$, $f'|f$.
- $C_{f,s} \subset C_{f',s'}$ if and only if $s'|s$, $f'|f$ and $f = \varphi_{s/s'}(x^{s'})f' \pmod{(1 - x)f'}$.

- $C_{f,s} \subset B_{f',s'}$ if and only if $s'|s$ and $f'|f(1-x)^{-1}$.
- $B_{f,s} \subset C_{f',s'}$ if and only if $s'|s$, $f'|f$ and s/s' is even.

In particular, the set of groups $B_{1-x^s,s}$, $s \geq 1$, is a basis for profinite topology.

Proof. By the Lemma 3.20 to any nontrivial normal subgroup V corresponds a triple (s, f, h) . If $s = 0$ then $V \subset \mathcal{A}_1$ and therefore V has infinite index. If $s \geq 1$ then the index of V is $sp^{\deg f}$ and by Definition 3.21 $V = B_{f,s}$ if $h = 0$ and $V = C_{f,s}$ if $h \neq 0$. Inclusions follow from the Lemma 3.1. To prove the last statement note that $B_{f,s} \supset B_{1-x^s,s}$, since $f|1-x^s$ by the definition of $B_{f,s}$. Also, $C_{f,s} \supset B_{1-x^{ps},ps}$, by the last item and the fact that $f|(1-x^s)^p = 1-x^{ps}$. Thus any normal subgroup of finite index contains some $B_{1-x^s,s}$, therefore it is a basis for profinite topology. \square

Now we describe the finite factors of \mathcal{L}_1 .

Theorem 3.23. *Any extension H of a finite cyclic group G by a finite elementary p -group E , such that the induced action of G on E has a cyclic vector, is a factor of \mathcal{L}_1 . In particular, if G has order s and m is a cyclic vector of E with minimal polynomial f (we normalize f so that $f(0) = 1$) then any kernel of a surjective homomorphism $\mathcal{L}_1 \rightarrow H$ is either $B_{f,s}$ or $C_{f,s}$. Moreover, if we fix action of G on E then there are at most two non-equivalent extensions. If $f(1) = 1$ or if $f(1) = 0$ but f does not divide $\varphi_s(x)$ then there is only split extension, in all other cases there are exactly two extensions.*

Proof. \mathcal{L}_1 can be given by $\{a, b | b^p = 1, [b^{a^n}, b^{a^m}] = 1 \text{ for all } n, m\}$. Therefore if H is an extension of G by E , G is generated by x and $m \in E$ is a cyclic vector, then the map $b \mapsto m$, $a \mapsto x$ can be lifted to a surjective map $\mathcal{L} \rightarrow H$.

The number of equivalent classes of extensions of G by E is equal to the order of $H^2(G, E)$, where $H^2(G, E)$ is the second cohomology group of G with coefficients in G -module E . By [7] $H^2(G, E) = E^G/NE$, where E^G is the set of fixed points in E under the action of G and $N = 1 + x + \dots + x^{s-1}$. If E has a cyclic vector m then it is isomorphic to $\mathbb{F}_p[x]/f\mathbb{F}_p[x]$ where f is the minimal polynomial for m . Thus if $g + (f)$ is a fixed point it follows that $xg = g \pmod{f}$, or equivalently $f|(1-x)g$. Since $\deg g < \deg f$ this can only happen if $f = (1-x)g$, i.e. if $f(1) = 0$ and $g = f(1-x)^{-1}$. Thus $\dim E^G = 1$ if $f(1) = 0$ and 0 otherwise. Also, $NE = \{1 + \dots + x^{s-1} + (f)\}$, and thus if $f(1) = 0$, $NE = E^G$ if and only if $f|(1 + \dots + x^{s-1})$. \square

3.4. Profinite and pro- p completion of $\mathcal{L}_{n,p}$. Perhaps the following lemma is well-known.

Lemma 3.24. *The profinite completion of \mathcal{L}_n is $\mathbb{F}_p[[\hat{\mathbb{Z}}]]^n \rtimes \hat{\mathbb{Z}}$, where $\hat{\mathbb{Z}}$ is the profinite completion of \mathbb{Z} and $\mathbb{F}_p[[\hat{\mathbb{Z}}]]$ is the profinite completion of the ring $R = \mathbb{F}_p[\mathbb{Z}]$, which is also the projective limit $\varprojlim \mathbb{F}_p[\mathbb{Z}/n\mathbb{Z}]$.*

Proof. First recall that any ideal of R is of the form fR where $f \in \mathbb{F}_p[x]$ and $f(0) = 1$. Now, any such irreducible f divides $1 - x^n$ for some n . Since f is the product of irreducibles it follows that for any f there is n such that $f \mid 1 - x^n$. Therefore the profinite completion of the ring $R = \mathbb{F}_p[x]$ is $\varprojlim R/(1 - x^n)R = \varprojlim \mathbb{F}_p[\mathbb{Z}/n\mathbb{Z}]$.

Consider now surjective maps $\psi_m : \mathbb{F}_p[[\hat{\mathbb{Z}}]]^n \rtimes \hat{\mathbb{Z}} \rightarrow \mathbb{F}_p[\mathbb{Z}/m\mathbb{Z}]^n \rtimes \mathbb{Z}/m\mathbb{Z} = \mathcal{L}_n/N_m$, where the normal subgroup N_m has triple $(m, (x - 1)^m \mathcal{A}_n, 0)$. The maps ψ_m are compatible with the projective system \mathcal{L}_n/N_m and therefore induce a map $\psi : \mathbb{F}_p[[\hat{\mathbb{Z}}]]^n \rtimes \hat{\mathbb{Z}} \rightarrow \varprojlim \mathcal{L}_n/N_m$. Then ψ is surjective by the Corollary 1.1.6 in [30], since $\mathbb{F}_p[[\hat{\mathbb{Z}}]]^n \rtimes \hat{\mathbb{Z}}$ is compact. On the other hand $\ker \psi_m = \ker \pi_m \times m\hat{\mathbb{Z}}$, where $\pi_m : \mathbb{F}_p[[\hat{\mathbb{Z}}]]^n \rightarrow \mathbb{F}_p[\mathbb{Z}/m\mathbb{Z}]^n$ is the natural projection. Since the intersection of $\ker \pi_m$ is trivial, the intersection of $\ker \psi_m$ is also trivial which implies that ψ is an isomorphism.

It is left to show that normal subgroups N_m form a basis of identity for the group \mathcal{L}_n , but it follows from Lemmas 3.1 and 3.2 that if V is normal subgroup of finite index in \mathcal{L}_n and $s > 0$ the generator of its projection on \mathbb{Z} then the normal subgroup with a triple $(sp, (x^{sp} - 1)\mathcal{A}_n, 0)$ lies in V . \square

Theorem 3.25. *Let V be the subgroup of \mathcal{L}_n with a triple (s, V_0, v) . Then it is profinitely closed in \mathcal{L}_n if and only if either $s > 0$ or $V = V_0$ is closed as a subset of $\mathbb{F}_p[[\hat{\mathbb{Z}}]]^n$ (that is, V is equal to the intersection of its closure with \mathcal{A}_n).*

Proof. If $s > 0$ then V is finitely generated by Theorem 3.5 and hence closed by Corollary 3.6. If $s = 0$ and $V = V_0 \subset \mathcal{A}_n$ which we identify with R^n , let \bar{V} denote the closure of V in $\mathbb{F}_p[[\hat{\mathbb{Z}}]]^n$. Then since $\mathbb{F}_p[[\hat{\mathbb{Z}}]]^n$ is compact \bar{V} is also compact and hence is still closed in $\mathbb{F}_p[[\hat{\mathbb{Z}}]]^n \rtimes \hat{\mathbb{Z}}$. Since $\mathbb{F}_p[[\hat{\mathbb{Z}}]]^n \cap \mathcal{L}_n = \mathbb{F}_p[[\hat{\mathbb{Z}}]]^n \cap R^n$, and hence $\bar{V} \cap \mathcal{L}_n = \bar{V} \cap R^n = V$. \square

Corollary 3.26. *There are subgroups of \mathcal{L}_1 which are not closed in the profinite topology.*

Proof. Here is an example due to D. Segal. Consider in $\mathcal{A}_1 = \sum_{\mathbb{Z}} (\mathbb{Z}/p\mathbb{Z})$ the subgroup H such that $v \in H$ if and only if $v_i = 0$ for i not a prime number. For any n the map $\mathbb{F}_p[\mathbb{Z}] \rightarrow \mathbb{F}_p[\mathbb{Z}/n\mathbb{Z}]$ maps H onto $\mathbb{F}_p[\mathbb{Z}/n\mathbb{Z}]$. Thus the profinite closure of H is $\mathbb{F}_p[[\hat{\mathbb{Z}}]]$, and so H is not closed (that is, the intersection of the closure of H with \mathcal{L}_1 is equal to $\mathcal{A}_1 \neq H$). \square

We will now study the pro- p completion of \mathcal{L}_1 . The index of $B_{f,s}$ or $C_{f,s}$ is equal to $p^{\deg f} s$, therefore we need to consider the cases when s is a power of p . In this case $1 - x^{p^n} = (1 - x)^{p^n}$, therefore $f = (1 - x)^i$ for $0 \leq i \leq p^n$. Thus we obtain a description of normal subgroups with index a power of p :

Definition 3.27. Define $B'_{i,n} = B_{(1-x)^i, p^n}$, $0 \leq i \leq p^n$ and $C'_{i,n} = C_{(1-x)^i, p^n}$, $1 \leq i \leq p^n$. We have that $B'_{i,n}$ and $C'_{i,n}$ is the list of all normal subgroups of \mathcal{L}_1 with index a power of p .

The following Theorem describes the lattice of normal subgroups of \mathcal{L}_1 of index a power of p .

Theorem 3.28. • $B'_{i,n} \subset B'_{j,m}$ if and only if $m \leq n$ and $j \leq i$.
 • $C'_{i,n} \subset C'_{j,m}$ if and only if either $m < n$, $j < i$ or $m = n$ and $j = i$.
 • $C'_{i,n} \subset B'_{j,m}$ if and only if $m \leq n$ and $j < i$.
 • $B'_{i,n} \subset C'_{j,m}$ if and only if $m < n$ and $j \leq i$.

In particular, the set of groups $B'_{p^n, n} = B_{1-x^{p^n}, p^n}$ is a basis for the pro- p topology.

Proof. Follows from Theorem 3.22 □

Corollary 3.29. The subgroup V of finite index in \mathcal{L}_1 is pro- p closed if and only if its triple (s, V_0, v) satisfies that $s = p^t \mathbb{Z}$ for some t and $V_0 \supset (1 - x^{p^n})\mathcal{A}_1$ for some $n \geq t$.

Lemma 3.30. The pro- p completion of \mathcal{L}_n is $\mathbb{F}_p[[\mathbb{Z}_p]]^n \rtimes \mathbb{Z}_p$, where \mathbb{Z}_p is the pro- p completion of \mathbb{Z} and $\mathbb{F}_p[[\mathbb{Z}_p]]$ is the pro- p completion of the ring $R = \mathbb{F}_p[\mathbb{Z}]$, which is also the projective limit $\varprojlim \mathbb{F}_p[\mathbb{Z}/p^n \mathbb{Z}]$.

Proof. The proof repeats almost verbatim the proof of Lemma 3.24, with $\hat{\mathbb{Z}}$ replaced by \mathbb{Z}_p . □

Lemma 3.31. Let V be a subgroup of \mathcal{L}_n with a triple (s, V_0, v) and $s > 0$. Then V is closed in the pro- p topology if and only if V_0 is equal to the intersection of its closure in $\mathbb{F}_p[[\mathbb{Z}_p]]^n$ with R^n , and for any $0 < q < s$ such that $|q|_p \geq |s|_p$

$$\frac{x^q - 1}{x^s - 1} v \notin \mathcal{A}_n + \bar{V}_0,$$

where \bar{V}_0 is the closure of V_0 in $\mathbb{F}_p[[\mathbb{Z}_p]]^n$ and $|q|_p$ is the highest power of p that divides q .

Proof. Let \bar{V} be the closure of V in $\mathbb{F}_p[[\mathbb{Z}_p]]^n \rtimes \mathbb{Z}_p$, and \bar{V}_0 the closure of V_0 in $\mathbb{F}_p[[\mathbb{Z}_p]]^n$. Then $\bar{V}_0 = \bar{V} \cap \mathbb{F}_p[[\mathbb{Z}_p]]^n$, hence if $V = \bar{V} \cap \mathcal{L}_n$ then

$$V_0 \subset \bar{V}_0 \cap \mathcal{A}_n \subset \bar{V} \cap \mathcal{A}_n \subset V \cap \mathcal{A}_n = V_0.$$

Suppose that V_0 is closed in $\mathbb{F}_p[[\mathbb{Z}_p]]^n$. Suppose that for any $0 < q < s$ such that $|q|_p \geq |s|_p$

$$\frac{x^q - 1}{x^s - 1}v \notin \mathcal{A}_n + \bar{V}_0.$$

Let $(w, c) \in \mathcal{L}_n$ be some element in \bar{V} . It means that there is a sequence $u_i \in V_0$ and $k_i \in \mathbb{Z}$ such that $(u_i + \varphi_{k_i}(x^s)v, k_i s)$ tends to (w, c) , or $(u_i + \varphi_{k_i}(x^s)v - x^{k_i s - c}w, k_i s - c)$ tends to identity. This happens if and only if

$$u_i + \varphi_{k_i}(x^s)v - x^{k_i s - c}w \rightarrow 0 \text{ in } \mathbb{F}_p[[\mathbb{Z}_p]]^n,$$

and $k_i s - c \rightarrow 0$ in \mathbb{Z}_p , which means that one can write $k_i s = p^{m_i}r_i + c$ and $m_i \rightarrow \infty$ as $i \rightarrow \infty$. It follows in particular that $|c|_p \geq |s|_p$. Note that $x^{k_i s - c} = x^{p^{m_i}r_i} \rightarrow 1$ in $\mathbb{F}_p[[\mathbb{Z}_p]] = \varprojlim \mathbb{F}_p[\mathbb{Z}/p^m\mathbb{Z}]$ since $x^{p^{m_i}r_i} = 1$ in $\mathbb{F}_p[\mathbb{Z}/p^m\mathbb{Z}]$ for $m \leq m_i$. For the same reason,

$$\varphi_{k_i}(x^s) = \frac{x^{k_i s} - 1}{x^s - 1} = \frac{x^{p^{m_i}r_i}x^c - 1}{x^s - 1} \rightarrow \frac{x^c - 1}{x^s - 1},$$

considered as a sequence in the field of fractions of $\mathbb{F}_p[[\mathbb{Z}_p]]$. It follows that u_i also has some limit $u \in \bar{V}_0$ and we have the equality

$$u + \frac{x^c - 1}{x^s - 1}v - w = 0.$$

Let $c = ks + q$ for $0 \leq q < s$. Since $|c|_p \geq |s|_p$ we have that $|q|_p \geq |s|_p$. We also have that $x^c - 1 = (x^s - 1 + 1)^k x^q - 1$, and therefore $x^c - 1 = g(x)(x^s - 1) + x^q - 1$ for some polynomial g . Thus

$$\frac{x^q - 1}{x^s - 1}v = \frac{x^c - 1}{x^s - 1}v - g(x)v = w - g(x)v - u \in R^n + \bar{V}_0,$$

which is by assumption impossible unless $q = 0$. Thus $c = ks$ and $w = u + \frac{x^{ks} - 1}{x^s - 1}v$, from which it follows that $u \in \bar{V}_0 \cap \mathcal{A}_n = V_0$ and therefore $(w, c) = (u, 0)(v, s)^k \in V$.

Conversely, suppose that there is $0 < q < s$ such that $|q|_p \geq |s|_p$ and

$$\frac{x^q - 1}{x^s - 1}v = w + u,$$

for $w \in \mathcal{A}_n$ and $u \in \bar{V}_0$. Choose a sequence $u_i \in V_0$ that tends to u , and a sequence $k_i \in \mathbb{Z}$ such that $k_i s \rightarrow q$ in \mathbb{Z}_p (it is possible to find since $|q|_p \geq |s|_p$). Then $(u_i, 0)(v, s)^{k_i} \in V$ but tend to (w, q) by reasoning similar to previous considerations. Hence V is not closed. \square

Corollary 3.32. $\mathbb{F}_p[[\mathbb{Z}_p]]$ is isomorphic to $\mathbb{F}_p[[t]]$, the ring of formal power series in t over \mathbb{F}_p . The map

$$a = \sum_{i \geq 0} a_i p^i \mapsto (1 + t)^a = \prod_{i \geq 0} (1 + t^{p^i})^{a_i},$$

where $a_i \in \{0, 1, \dots, p-1\}$, is a well-defined isomorphism of \mathbb{Z}_p into the multiplicative group of $\mathbb{F}_p[[t]]$ and the action of \mathbb{Z}_p on $\mathbb{F}_p[[t]]$ in $\mathbb{F}_p[[\mathbb{Z}_p]]^n \rtimes \mathbb{Z}_p$ ($\mathbb{F}_p[[\mathbb{Z}_p]]$ identified with $\mathbb{F}_p[[t]]$) in Lemma 3.30 is given by the formula $a * f(t) = (1+t)^a f(t)$.

Proof. Note that any element of $\mathbb{F}_p[[\mathbb{Z}_p]] = \varprojlim R/(1-x^{p^n})R = \varprojlim R/(1-x)^{p^n}R = \varprojlim_j R/(1-x)^j R$ can be written as $\sum_{i \geq 0} \varepsilon_i (x-1)^i$, for $\varepsilon_i \in \mathbb{F}_p$. Then the map $\sum_{i \geq 0} \varepsilon_i (x-1)^i \mapsto \sum_{i \geq 0} \varepsilon_i t^i$ gives an isomorphism of rings $\varprojlim R/(1-x)^i R$ and $\mathbb{F}_p[[t]]$. The other claims are obvious. \square

Lemma 3.33. *Let $g = (x-1)^c h$ for some $g, h \in R$, where $x-1$ does not divide h . Then $g\mathbb{F}_p[[\mathbb{Z}_p]] \cap R = (x-1)^c R$.*

Proof. Note that multiplying by some positive power of x we may assume that $g, h \in \mathbb{F}_p[x]$. Hence it suffices to prove that if $g \in \mathbb{F}_p[x]$, we have that $g\mathbb{F}_p[[\mathbb{Z}_p]] \cap \mathbb{F}_p[x] = (x-1)^c \mathbb{F}_p[x]$. Change variable $t = x-1$ and note that $\mathbb{F}_p[x] = \mathbb{F}_p[t]$. Now, using first part of Corollary 3.32 we have to prove that if $g \in \mathbb{F}_p[t]$, $g = t^c h$ and $h(0) \neq 0$ then $g\mathbb{F}_p[[t]] \cap t^c \mathbb{F}_p[t] = t^c \mathbb{F}_p[t]$. Notice that since $h(0) \neq 0$, h is invertible in $\mathbb{F}_p[[t]]$ and therefore $g\mathbb{F}_p[[t]] = t^c \mathbb{F}_p[[t]]$. It is left to recall that the element $\sum_{j \geq 0} \varepsilon_j t^j \in \mathbb{F}_p[[t]]$ belongs to $\mathbb{F}_p[t]$ if and only if the number of nonzero ε_j is finite. \square

Lemma 3.34. *Let $U \subset R^n$ be a subgroup such that $e(U) = p^e$ for some $e \geq 0$. Let $k = np^e$. Then there are elements $h_1, \dots, h_k \in R^n$ and $d_1, \dots, d_k \in \mathbb{F}_p[x]$ such that $d_j = 0$ or $d_j(0) = 1$, $R^n = \bigoplus_{j=1}^k R^{p^e} h_j$ and $U = \bigoplus_{j=1}^k (Rd_j)^{p^e} h_j$. Moreover, U is equal to the intersection of its closure in $\mathbb{F}_p[[\mathbb{Z}_p]]^n$ with R^n if and only if $\det^* U = \prod_{j: d_j \neq 0} d_j$ is a power of $1-x$ (recall the definitions 2.9 and 2.7).*

Proof. The existence of such h_j and d_j follows using the fact that R^n seen as R module with the action $x*v = x^{p^e}v$ is isomorphic to R^k (to see this, consider a formula similar to (2.1)) and then applying Lemma 2.3.

To prove the second part note first that $\mathbb{F}_p[[\mathbb{Z}_p]]^n = \bigoplus_{j=1}^k (\mathbb{F}_p[[\mathbb{Z}_p]])^{p^e} h_j$. Indeed, suppose that $u_j \in \mathbb{F}_p[[\mathbb{Z}_p]]$ and $\sum_j u_j^{p^e} h_j = 0$. Let $u_j = \lim_q u_{j,q}$ so that $u_{j,q} \in R$ and $u_j - u_{j,q} \in (x-1)^q \mathbb{F}_p[[\mathbb{Z}_p]]$. It follows that $\sum_j u_{j,q}^{p^e} h_j \in (x-1)^q \mathbb{F}_p[[\mathbb{Z}_p]] \cap R^n = (x-1)^q R^n$ by Lemma 3.33. Hence $\sum_j u_{j,q}^{p^e} h_j = (x-1)^q \sum_j v_{j,q}^{p^e} h_j$ for some $v_{j,q} \in R$. Thus for $q > p^e$ we obtain $\sum_j (u_{j,q} - (x-1)^{q-p^e} v_{j,q})^{p^e} h_j = 0$. It follows that $u_{j,q} = (x-1)^{q-p^e} v_{j,q}$ and therefore $u_j = \lim_q u_{j,q} = 0$.

It follows that if \bar{U} is the closure of U in $\mathbb{F}_p[[\mathbb{Z}_p]]^n$, then

$$\bar{U} = \bigoplus_{j=1}^k (\mathbb{F}_p[[\mathbb{Z}_p]])^{p^e} d_j h_j.$$

Let $d_j = (x - 1)^{r_j} h_j$ for some $r_j \geq 0$ and h_j such that $x - 1$ does not divide h_j . Then by Lemma 3.33 we obtain that

$$\begin{aligned} \bar{U} \cap R^n &= \oplus_{j=1}^k (\mathbb{F}_p[[\mathbb{Z}_p]] d_j)^{p^e} h_j \cap R^n = \\ &= \oplus_{j=1}^k (\mathbb{F}_p[[\mathbb{Z}_p]] (x - 1)^{r_j})^{p^e} h_j \cap R^n = \oplus_{j=1}^k (R(x - 1)^{r_j})^{p^e} h_j, \end{aligned}$$

hence $U = \bar{U} \cap R^n$ if and only if $h_j = 1$. \square

Theorem 3.35. *Let V be a subgroup of \mathcal{L}_n with a triple (s, V_0, v) . Then V is closed in the pro- p topology if and only if either $s = 0$ and V_0 is closed as a subset of $\mathbb{F}_p[[\mathbb{Z}_p]]^n$, or $s > 0$ and for V_0 we have that $e(V_0)$ divides $p^{|s|_p}$, $\det^* V_0$ is a power of $1 - x$, and for any $0 < q < s$ such that $|q|_p \geq |s|_p$*

$$(x^q - 1)v \notin (x^s - 1)\mathcal{A}_n + (x - 1)^{p^{|s|_p}} V_0.$$

Proof. The case $s = 0$ is obvious, so assume $s > 0$. If V is pro- p closed then it follows that V is the intersection of the p -chain V_j with triples (s_j, V_j, v_j) . Since it is p -chain, each s_j is a power of p , and since $s_j | s$, the sequence s_j stabilize. Thus eventually $s_j = p^{e'}$ for some $e' \leq e$. Then eventually $x^{p^{e'}} V_j = V_j$, and since $V_0 = \cap V_j$, we obtain that also $x^{p^e} V_0 = V_0$. Then since V is pro- p closed, V_0 is closed in $\mathbb{F}_p[[\mathbb{Z}_p]]^n$ and therefore by Lemma 3.34 $\det^* V_0$ is a power of $1 - x$.

Conversely, suppose that $x^{p^e} V_0 = V_0$ and $\det^* V_0$ is a power of $1 - x$. By Lemma 3.34 V_0 is closed in $\mathbb{F}_p[[\mathbb{Z}_p]]^n$.

Notice also that

$$\frac{x^q - 1}{x^s - 1} v \notin R^n + \bar{V}_0$$

if and only if $(x^q - 1)v \notin (x^s - 1)R^n + ((x^s - 1)\bar{V}_0) \cap R^n$. Note that we can write $x^s - 1 = ((x - 1)h')^{p^e}$ where h' is not divisible by $x - 1$, and therefore invertible in $\mathbb{F}_p[[\mathbb{Z}_p]]$. Using the decomposition from the Lemma 3.34 we have that $\bar{V}_0 = \oplus_{j=1}^k (\mathbb{F}_p[[\mathbb{Z}_p]] d_j)^{p^e} h_j$, hence

$$(x^s - 1)\bar{V}_0 \cap R^n = \oplus_{j=1}^k (\mathbb{F}_p[[\mathbb{Z}_p]] (x - 1) d_j)^{p^e} h_j \cap R^n = \oplus_{j=1}^k (R(x - 1) d_j)^{p^e} h_j.$$

It follows that

$$((x^s - 1)\bar{V}_0) \cap R^n = \oplus_{j=1}^k (R(x - 1) d_j)^{p^e} h_j = (x - 1)^{p^e} V_0.$$

\square

3.5. Automaton that generates $\mathcal{L}_{1,p}$. In this subsection we give description of a part of lattice of subgroups in $\mathcal{L}_{1,p}$ in terms of action on a p -regular rooted tree. We also provide automaton presentation of $\mathcal{L}_{1,p}$ showing, in particular, that groups $\mathcal{L}_{1,p}$ are self-similar.

Consider in the group $\mathcal{L}_1 = \mathcal{L}_{1,p}$ the p -chain $H_m = (1 - x)^m R \rtimes \mathbb{Z}$, for $m \geq 0$, that is H_m has the triple $(1, (1 - x)^m R, 0)$. Since this is p -chain the index of H_m is p^m . By the Lemma 3.2 each H_m is normal, in

particular the chain is subnormal. Identify the p -regular rooted tree T_p with the set of words over the alphabet $\Omega = \{0, 1, \dots, p-1\}$. For each $\omega \in \Omega^*$, $\omega = \omega_0 \cdots \omega_{m-1}$ construct the element $\bar{\omega} = \sum_{i=0}^{m-1} \omega_i (x-1)^i \in R$. The following Lemma holds:

Lemma 3.36. $\omega \mapsto (\bar{\omega}, 0)H_{|\omega|}$ is the isomorphism of T_p and the coset tree of the chain $\{H_m\}$, where $|\omega|$ is the length of the word ω .

Proof. For injectivity, consider that $(\bar{\omega}, 0)H_{|\omega|} = (\bar{\theta}, 0)H_{|\theta|}$ if and only if $|\theta| = |\omega| = m$ and $\bar{\theta} - \bar{\omega} \in (1-x)^m R$, that is if and only if $\bar{\omega} = \bar{\theta}$, which means that $\omega_i = \theta_i$ and hence $\omega = \theta$. For surjectivity, consider that for any $(w, s) \in \mathcal{L}_1$ we have that $(w, s)H_m = (w, 0)H_m$. Note also that $R/(1-x)^m R = \mathbb{F}_p[x]/\langle(1-x)^m\rangle$, so for any $w \in R$ there is a sequence $w_i \in \Omega$, such that

$$w = \sum_{i=0}^{m-1} w_i (x-1)^i \pmod{(1-x)^m R},$$

for any $m \geq 0$. Finally, it is easy to establish that the map $\omega \mapsto \bar{\omega}$ preserve the property of vertices being adjoint. \square

In the following Proposition we present the description of the stabilizers of the action of \mathcal{L}_1 on the coset tree of $\{H_m\}$, identified with T_p with the help of the map from the above lemma:

Proposition 3.37. *The subgroup $\text{Stab}_{\mathcal{L}_1}(\omega)$ has the triple $(1, (1-x)^{|\omega|}R, (1-x)\bar{\omega})$, and the stabilizer of the m -th level $\text{Stab}_{\mathcal{L}_1}(m)$ has the triple $(p^e, (1-x)^m R, 0)$, where e is smallest such that $m \leq p^e$.*

Proof. Note that in the action on the coset tree the stabilizer of the vertex gH_m is $gH_m g^{-1}$. Hence $\text{Stab}_{\mathcal{L}_1}(\omega) = (\bar{\omega}, 0)H_{|\omega|}(-\bar{\omega}, 0)$, from which the formula for the triple follows.

To find $\text{Stab}_{\mathcal{L}_1}(m) = \cap_{\omega: |\omega|=m} \text{Stab}_{\mathcal{L}_1}(\omega)$ use Corollary 3.4. It follows that $\text{Stab}_{\mathcal{L}_1}(m)$ has the triple $(r, (1-x)^m R, v)$ where $r > 0$ is minimal such that $(1-x^r)\bar{\omega} = (1-x^r)\bar{\omega}' \pmod{(1-x)^m R}$ for any ω, ω' of length m , and $v = (1-x^r)\bar{\omega}$. Choosing $\omega = 0^m$ we notice that we can take $v = 0$ and $r > 0$ is minimal such that $(1-x^r)\bar{\omega}' = 0 \pmod{(1-x)^m R}$ for all ω' of length m . Equivalently, $r > 0$ is minimal such that $(1-x)^m$ divides $1-x^r$, that is $x^r = 1 + (x-1)^m f(x)$ for some $f \in \mathbb{F}_p[x]$. Let $r = p^e r'$ and change variable x to $y = x-1$, then we have that $\cdots + r' y^{p^e} + 1 = (y^{p^e} + 1)^{r'} = (y+1)^r = 1 + y^m f(y+1)$, and so since $r' \neq 0$ in \mathbb{F}_p one must have that $p^e \geq m$, hence the minimal such $r = p^e$ with $e \geq \log_p(m)$. \square

Now we will describe the stabilizers of infinite paths. To do this, we need a little preparation. By the proof of the Corollary 3.32 $\varprojlim_j R/(1-x)^{p^j}$

$x)^m R = \mathbb{F}_p[[t]]$, and the subalgebra of $\mathbb{F}_p[[t]]$ generated by t and $(t-1)^{-1} = -\sum_{i \geq 0} t^i$ is isomorphic to R by the isomorphism that maps $t-1$ to x , so we may consider R as a subalgebra of $\mathbb{F}_p[[t]]$. Then any $\omega \in \Omega^{\mathbb{N}}$ defines an element $\bar{\omega} \in \mathbb{F}_p[[t]]$ by the rule $\bar{\omega} = \sum_{i \geq 0} \omega_i t^i$. Let $\mathbb{F}_p((t))$ be the ring of fractions of $\mathbb{F}_p[[t]]$, then we can consider $(1-x^s)^{-1}R$ as a subset of it.

Proposition 3.38. *For $\omega \in \Omega^{\mathbb{N}}$ the $\text{Stab}_{\mathcal{L}_1}(\omega)$ is the cyclic group generated by (w, s) in case $s > 0$ is the smallest integer such that $\bar{\omega} \in (1-x^s)^{-1}R$, and $w = (1-x^s)\bar{\omega}$. In case there is no such s the $\text{Stab}_{\mathcal{L}_1}(\omega)$ is trivial.*

Proof. Denote by $\omega^{(m)}$ prefix of length m of ω . Note that $\cap_m (1-x)^m R$ is trivial, so $\text{Stab}_{\mathcal{L}_1}(\omega) = \cap_m \text{Stab}_{\mathcal{L}_1}(\omega^{(m)})$ can be either trivial or generated by some (w, s) for $s > 0$. The second case happens if and only if for every m we have $(w, s) \in \text{Stab}_{\mathcal{L}_1}(\omega^{(m)})$, and s is smallest that this is possible. Since by Proposition 3.37 the triple of $\text{Stab}_{\mathcal{L}_1}(\omega^{(m)})$ is $(1, (1-x)^m R, \sum_{i=0}^{m-1} \omega_i (x-1)^{i+1})$, and

$$\left(\sum_{i=0}^{m-1} \omega_i (x-1)^{i+1}, 1\right)^s = ((1-x^s) \left(\sum_{i=0}^{m-1} \omega_i (x-1)^i\right), s),$$

this can happen if and only if $w = (1-x^s) \left(\sum_{i=0}^{m-1} \omega_i (x-1)^i\right) \pmod{(1-x)^m R}$ for every m , or equivalently that $w - (1-x^s)\bar{\omega} \in (1-x)^m R$. Since $\cap_m (1-x)^m R$ is trivial, it means that $w = (1-x^s)\bar{\omega}$. \square

We finally describe the automaton that generates the action of $\mathcal{L}_{1,p}$ on the tree.


Proposition 3.39. *Let $b = (e, 0)$ and $a = (0, 1)$ be the generators of \mathcal{L}_1 . Then their action on vertices of the tree can be described as*

$$(3.3) \quad \begin{aligned} b(\omega_0 \omega_1 \dots \omega_m) &= (\omega_0 + 1) \omega_1 \dots \omega_m, \\ a(\omega_0 \omega_1 \dots \omega_m) &= \omega_0 (\omega_1 + \omega_0) \dots (\omega_m + \omega_{m-1}), \end{aligned}$$


where the addition is mod p . The relations 3.3 lead to the following recursive formula $a = (a, ba, \dots, b^{p-1}a)$ describing the action of a in terms of its projections on subtrees with roots at first level. Denoting $a_i = b^i a$, $0 \leq i \leq p-1$, we see that \mathcal{L}_1 is generated by the automaton with p states a_0, \dots, a_{p-1} .

Remark 5. *Note that for $p = 2$ it is the automaton studied in [19].*

Here are Moore diagrams of automata generating \mathcal{L}_p for $p = 2$:



and $p = 3$:



4. APPLICATION TO RELATIVE GRADIENT RANK

As was already mentioned in the introduction, Lackenby defined an interesting group theoretical notion, the rank gradient of a group, which happens to be useful in topology, studies on orbit equivalence, on amenability properties of groups and other topics [25]. Given a group G and a descending sequence $\{H_i\}_{i=1}^{\infty}$ of subgroups of finite index the rank gradient of the sequence $\{H_i\}$ with respect to G is defined as

$$RG(G, \{H_i\}) = \lim_{i \rightarrow \infty} \frac{d(H_i) - 1}{[G : H_i]},$$

where $d(H)$ denotes the minimal number of generators of a group H .

The notion of amenable groups was introduced by J. von Neumann in 1929 and plays important role in many areas of mathematics [8]. There are a number of results due to Lackenby, Abert, Jaikin-Zapirain, Luck and Nikolov, showing that amenability of G or of a normal subgroups of G implies vanishing of the rank gradient. For instance, finitely generated infinite amenable groups have zero gradient rank with respect to any normal chain with trivial intersection (see Theorem 5 in [1]). Obviously it is reasonable to restrict study of the rank gradient for sequences $\{H_i\}$ with trivial core (i.e. no nontrivial normal subgroups in the intersection $\cap_i H_i$). The attention was mostly to the case when $\cap_i H_i = \{1\}$ but the case $\cap_i H_i \neq \{1\}$ is also interesting.

Question 1. [1] *Let G be a finitely generated infinite amenable group. Is it true that $RG(G, \{H_i\}) = 0$ for any chain with trivial intersection? One can also ask whether the following stronger statement is true: Is it true that $RG(G, \{H_i\}) = 0$ if the chain has trivial core?*

In case of groups \mathcal{L}_n the answer to Question 1 is positive:

Theorem 4.1. *Let H_i , $i \geq 1$ be a descending chain of subgroups in \mathcal{L}_n with trivial core, then $RG(\mathcal{L}_n, \{H_i\}) = 0$. If $n = 1$ the converse is also true.*

Proof. Let (s_i, V_i, v_i) be the triples of H_i , and let $m_i = \dim_{\mathbb{F}_p} \mathcal{A}_n / V_i$. By the Theorem 3.5 H_i is isomorphic to \mathcal{L}_{ns_i} , thus $d(H_i) = ns_i + 1$. On the other hand $[\mathcal{L}_n : H_i] = s_i p^{m_i}$, thus $RG(\mathcal{L}_n, \{H_i\}) = 0$ if and only if $m_i \rightarrow \infty$ when $i \rightarrow \infty$. Suppose $m_i \not\rightarrow \infty$. Since the chain is descending, m_i is nondecreasing, therefore the sequence $\{m_i\}$ stabilizes, that is, there is such i_0 that $V_{i_0} = V_j$ for $j \geq i_0$, and let us denote it $V = V_{i_0}$. Let $W = \cap_{t \in \mathbb{Z}} x^t V$, then $xW = W$ and thus W is a normal subgroup of \mathcal{L}_n that is contained in all H_i , $i \geq 1$. Note that W actually equals to the finite intersection $\cap_{t=1}^{t=s_0} x^t V$ since by the Lemma 3.2 $x^{s_0} V = V$. Therefore W has finite index in \mathcal{A}_n and thus is not trivial. It follows that the core of $\{H_i\}$ is also not trivial, since it contains W .

Let $n = 1$. Above we showed that $RG(\mathcal{L}_n, \{H_i\}) \neq 0$ if and only if $V_i = H_i \cap \mathcal{A}_n$ stabilizes. Let N be the core of $\{H_i\}$ and suppose N is not trivial. Then it follows from Corollary 3.3 that $N' = N \cap \mathcal{A}$ has finite index in \mathcal{A}_1 . Since N' is contained in each V_i , the chain $\{V_i\}$ must stabilize. \square

If $\cap_{i=1}^{\infty} H_i = H$ then H is a closed subgroup with respect to the profinite topology and $RG(G, \{H_i\})$ is a characteristic of the pair (G, H) which in some situations may characterize the pair (G, H) up to isomorphism (two pairs $(G, H), (P, Q)$ are isomorphic if there is an isomorphism $\phi : G \rightarrow P$ such that $\phi(H) = Q$). If $RG(G, \{H_i\}) = 0$ then the function

$$rg(m) = rg_{(G, \{H_i\})}(m) = \frac{d(H_m) - 1}{[G : H_m]},$$

which we call *relative rank gradient (function)* can be a subject of investigation (we omit the index $(G, \{H_i\})$ if the group and chain in consideration are understood). The rate of decay of $rg(m)$ is a sort of characteristic of the pair (G, H) and may characterize the way H is embedded in G as a subgroup. Note that the same subgroup can be obtained as intersection of distinct chains. For instance, if the chain consists of subgroups isomorphic to the enveloping group (and such chains exist in scale invariant groups for instance) one can delete certain elements in H_i thereby allowing $rg(m)$ to decay as fast as one would

like. Therefore, it is reasonable to restrict attention to the case when for some prime p the index $[H_i : H_{i+1}] = p$. In this case we say chain is a p -chain. One of corollaries of our results listed above is showing that a lamplighter group contains subnormal p -chains of trivial intersection, with distinct rates of decay of the rank gradient function.

Let H_i , $i \geq 0$ be a descending p -chain of subgroups of \mathcal{L}_n such that $H_0 = \mathcal{L}_n$. Note that for each i the subgroup H_i is isomorphic to \mathcal{L}_k for some k by Theorem 3.5. Then there are two possible cases for H_{i+1} : either its projection on \mathbb{Z} (under the identification of H_i and \mathcal{L}_k) is \mathbb{Z} or it is $p\mathbb{Z}$. In the first case $d(H_{i+1}) = d(H_i)$ and in the second case $d(H_{i+1}) = pd(H_i) - p + 1$. We have the following result

Theorem 4.2. *Suppose that $g : \mathbb{N} \rightarrow \mathbb{N}$ is such that $g(0) = p + 1$ and for each $i \in \mathbb{N}$ either $g(i + 1) = g(i)$ or $g(i + 1) = pg(i) - p + 1$, and suppose that the set $\{i | g(i) = g(i + 1)\}$ is infinite. Then there is a descending subnormal p -chain $\{H_i\}$ of subgroups of \mathcal{L}_1 , such that $H_0 = \mathcal{L}_1$, $d(H_i) = g(i)$ and $\cap_i H_i = \{1\}$.*

Proof. Note that by Theorem 3.11 we can construct some descending subnormal p -chain $\{H'_i\}$ such that all of the above properties hold except possibly for $\cap_i H'_i = \{1\}$. Note also that

$$\begin{aligned} \frac{g(i+1) - 1}{p^{i+1}} &= \frac{g(i) - 1}{p^i} \text{ if } g(i+1) = pg(i) - p + 1, \\ \frac{g(i+1) - 1}{p^{i+1}} &= \frac{1}{p} \frac{g(i) - 1}{p^i} \text{ if } g(i+1) = g(i). \end{aligned}$$

Since $\{H'_i\}$ is a p -chain, $[\mathcal{L}_1 : H_i] = p^i$, and the fact that the set $\{i | g(i+1) = g(i)\}$ is infinite implies that $RG(\mathcal{L}_1, \{H'_i\}) = 0$. By the Theorem 4.1 it follows that the core of the chain $\{H'_i\}$ is trivial.

Consider then the rooted binary tree associated to this chain, and the action of G on it by multiplication from the right. The kernel of this action is the core of the chain $\{H'_i\}$, and therefore the action is faithful. It is clear that it is also spherically transitive. Then it follows from the Corollary 3.15 and the Proposition 4.11 in [15] that the action of \mathcal{L}_1 on the boundary of this tree is topologically free, thus the set of infinite paths in the tree with nontrivial stabilizer is a union of nowhere dense subsets of the boundary of the tree. In particular its complement is not empty. Take a point from this complement, that is an infinite path $\{g_i H'_i\}$ (for some $g_i \in \mathcal{L}_1$) with trivial stabilizer. It is easy to see that this stabilizer is the intersection of descending subnormal p -chain $H_i = g_i H'_i g_i^{-1}$. To finish the proof it is left to note that since H'_i and H_i are isomorphic, $d(H_i) = d(H'_i) = g(i)$ for all $i \geq 0$. \square

Being solvable groups \mathcal{L}_n are elementary amenable groups. The p -groups (p prime) of intermediate growth constructed in [13] are nonelementary amenable groups. Since for them gradient rank is also zero it is reasonable to study relative gradient rank for various closed subgroups in pro- p -topology and p -chains representing them.

One of such groups is the group $\mathcal{G} = \langle a, b, c, d \rangle$ generated by 4 involutions [20]. This group (as well as most of other known groups of intermediate growth) act faithfully by automorphisms on a regular rooted tree (in this case on a binary tree). We identify in a standard way vertices of binary tree with binary words. For the stabilizer $P = st_{\mathcal{G}}(1^\infty)$ of infinite ray 1^∞ (point of the boundary ∂T of a tree) the corresponding sequence is the sequence $P_m = st_{\mathcal{G}}(1^m)$. As is shown in [], $P_1 = st_{\mathcal{G}}(1) = \langle c, d, c^a, d^a \rangle$, $P_m = R_m \rtimes \{1, b, c, d\}$ for $m \geq 2$, where R_m are iterative semidirect products $R_1 = K = \langle (ab)^2 \rangle^{\mathcal{G}}$, and $R_m = (K \times R_{m-1}) \rtimes \{1, (ac)^4\}$ for $m \geq 2$. Computations then show that $d(P_m) = m + 4$ for $m \geq 2$, $d(P_1) = 4$. This implies that

$$rg(\mathcal{G}, \{P_i\})(m) = \frac{m + 4}{2^m},$$

for $m \geq 2$.

Question 2. Which subgroups of \mathcal{G} are closed in profinite topology and what kind of asymptotic behavior $rg(m)$ holds for them?

REFERENCES

- [1] Miklós Abért, Andrei Jaikin-Zapirain, and Nikolay Nikolov. The rank gradient from a combinatorial viewpoint. *Groups Geom. Dyn.*, 5(2):213–230, 2011.
- [2] D. J. Allums and R. I. Grigorchuk. The rank gradient and the lamplighter group. *to appear in Involve Journal of Mathematics*.
- [3] M. F. Atiyah. Elliptic operators, discrete groups and von Neumann algebras. In *Colloque “Analyse et Topologie” en l’Honneur de Henri Cartan (Orsay, 1974)*, pages 43–72. Astérisque, No. 32–33. Soc. Math. France, Paris, 1976.
- [4] Tim Austin. Rational group ring elements with kernels having irrational dimension. 2009.
- [5] C. Bendikov, A. Pittet and R. Sauer. Spectral distribution and l^2 -isoperimetric profile of laplace operators on groups. 2009.
- [6] Ievgen Bondarenko, Rostislav Grigorchuk, Rostyslav Kravchenko, Yevgen Muntyan, Volodymyr Nekrashevych, Dmytro Savchuk, and Zoran Šunić. Groups generated by 3-state automata over a 2-letter alphabet. I. *São Paulo J. Math. Sci.*, 1(1):1–39, 2007.
- [7] Kenneth S. Brown. *Cohomology of groups*, volume 87 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1994. Corrected reprint of the 1982 original.
- [8] Tullio Ceccherini-Silberstein, Rostislav Grigorchuk, and Pierre de la Harpe. Décompositions paradoxales des groupes de Burnside. *C. R. Acad. Sci. Paris Sér. I Math.*, 327(2):127–132, 1998.

- [9] Mahlon M. Day. Amenable semigroups. *Illinois J. Math.*, 1:509–544, 1957.
- [10] Yves de Cornulier. Finitely presented wreath products and double coset decompositions. *Geom. Dedicata*, 122:89–108, 2006.
- [11] Anna Erschler. Generalized wreath products. *Int. Math. Res. Not.*, pages Art. ID 57835, 14, 2006.
- [12] Łukasz Grabowski. On the Atiyah problem for the lamplighter groups. 2010.
- [13] R. I. Grigorchuk. Construction of p -groups of intermediate growth that have a continuum of factor-groups. *Algebra i Logika*, 23(4):383–394, 478, 1984.
- [14] R. I. Grigorchuk. Just infinite branch groups. In *New horizons in pro- p groups*, volume 184 of *Progr. Math.*, pages 121–179. Birkhäuser Boston, Boston, MA, 2000.
- [15] R. I. Grigorchuk. Some topics in the dynamics of group actions on rooted trees. *Proc. Steklov Inst. Math.*, 273:72–191, 2011.
- [16] R. I. Grigorchuk, T. Chekerini-Sil’berstaĭn, and P. de la Harpe. Amenability and paradoxical decompositions for pseudogroups and discrete metric spaces. *Tr. Mat. Inst. Steklova*, 224(Algebra. Topol. Differ. Uravn. i ikh Prilozh.):68–111, 1999.
- [17] R. I. Grigorchuk, V. V. Nekrashevich, and V. I. Sushchanskiĭ. Automata, dynamical systems, and groups. *Tr. Mat. Inst. Steklova*, 231(Din. Sist., Avtom. i Beskon. Gruppy):134–214, 2000.
- [18] Rostislav I. Grigorchuk, Peter Linnell, Thomas Schick, and Andrzej Żuk. On a question of Atiyah. *C. R. Acad. Sci. Paris Sér. I Math.*, 331(9):663–668, 2000.
- [19] Rostislav I. Grigorchuk and Andrzej Żuk. The lamplighter group as a group generated by a 2-state automaton, and its spectrum. *Geom. Dedicata*, 87(1-3):209–244, 2001.
- [20] R. I. Grigorčuk. On Burnside’s problem on periodic groups. *Funktsional. Anal. i Prilozhen.*, 14(1):53–54, 1980.
- [21] P. Hall. On the finiteness of certain soluble groups. *Proc. London Math. Soc.* (3), 9:595–622, 1959.
- [22] Thomas W. Hungerford. *Algebra*, volume 73 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1980.
- [23] V. A. Kaĭmanovich and A. M. Vershik. Random walks on discrete groups: boundary and entropy. *Ann. Probab.*, 11(3):457–490, 1983.
- [24] P. H. Kropholler. A note on the cohomology of metabelian groups. *Math. Proc. Cambridge Philos. Soc.*, 98(3):437–445, 1985.
- [25] Marc Lackenby. Expanders, rank and graphs of groups. *Israel J. Math.*, 146:357–370, 2005.
- [26] Franz Lehner and Stephan Wagner. Free lamplighter groups and a question of atiyah. 2010.
- [27] Alexander Lubotzky and Dan Segal. *Subgroup growth*, volume 212 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 2003.
- [28] Volodymyr Nekrashevych and Gábor Pete. Scale-invariant groups. *Groups Geom. Dyn.*, 5(1):139–167, 2011.
- [29] Schick Thomas Pichot, Mikaël and Andrzej Żuk. Closed manifolds with transcendental L2-Betti numbers. 2010.
- [30] Luis Ribes and Pavel Zalesskii. *Profinite groups*, volume 40 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series*

of Modern Surveys in Mathematics]. Springer-Verlag, Berlin, second edition, 2010.

- [31] J. B. Riles. The near Frattini subgroups of infinite groups. *J. Algebra*, 12:155–171, 1969.

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843, USA

E-mail address: grigorch@math.tamu.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, CHICAGO, IL 60637, USA

E-mail address: rkchenko@math.uchicago.edu